

SACOX® 120 MicroGranulate
Premix for medicated feeding stuff
 Keep out of reach of children.

STATEMENT OF ACTIVE SUBSTANCE

Salinomycin sodium 120 mg

PRODUCT DESCRIPTION

SACOX® 120 MicroGranulate represents a beige to brown dust free granular product for incorporation into animal feedingstuffs.

PHARMACODYNAMICS

Salinomycin sodium is active against certain Gram-positive bacteria, while Gram-negative species are resistant. The use of salinomycin sodium as a feed additive at the proposed concentration is unlikely to increase shedding of *Salmonella*, *E. coli* and *Campylobacter* or to induce resistance and cross-resistance to antimicrobials important in human and animal therapy.

PHARMACOKINETICS

Salinomycin is absorbed to a certain extent in the chicken and extensively metabolised. Unchanged salinomycin represents a very small fraction of the metabolites excreted.

Many metabolites, predominantly mono- and multi-hydroxylated, have been identified in the excreta and in the tissues. The metabolites in excreta show a higher degree of hydroxylation than in the liver. Salinomycin-related metabolites have a reduced ionophoric activity when compared to salinomycin.

Salinomycin is considered as the marker residue. No residues in eggs are expected provided that the proposed maximum dose and duration of administration are respected.

INDICATIONS FOR USE

For the prevention of coccidiosis caused by *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. mivati*, *E. necatrix* and *E. tenella* in broiler chickens and in replacement birds intended for use as caged layers.

RECOMMENDED DOSAGE:

POULTRY: Thoroughly mix 500 g of SACOX per one tonne of finished feed. This is equivalent to 60 g of Salinomycin sodium per tonne of feed (60 ppm), which should be fed continuously as the sole ration.

MODE OF ADMINISTRATION

For oral administration in feed.

To assist in correct mixing the required amount of SACOX should be pre-mixed with at least 9.5 kg ground meal, or the normal vitamin and mineral premix, before gradually adding to the bulk feed.

CONTRAINDICATIONS

The recommended Inclusion dose rates should not be exceeded.

This product must not be fed to horses or any other animal or bird not included on the label claim.

WARNINGS AND PRECAUTIONS

For animal treatment only.

Harmful if swallowed. May irritate the eyes. Avoid contact with the eyes. Do not inhale dust. When opening the container and mixing into feed, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length PVC gloves, goggles

and disposable dust face mask covering mouth and nose. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, goggles and contaminated clothing.

INTERACTIONS WITH OTHER MEDICAMENTS

Other anticoccidial drugs should not be added to feeds containing SACOX.

SACOX should not be used concurrently with Tiamulin or other antibiotics.

STATEMENT ON USAGE DURING PREGNANCY AND LACTATION

DO NOT USE in birds laying eggs for human consumption or within 7 days of laying where eggs or egg products are to be used for human consumption or processing.

ADVERSE EFFECTS/UNDESIRABLE EFFECTS

For chickens reared for laying, 50 mg Salinomycin sodium per kg complete feed is considered safe for a feeding period of the first 12 weeks of life.

OVERDOSE AND TREATMENT

Salinomycin sodium levels up to 120 mg/kg feed are tolerated for 35 days without effects on any of the end-points measured.

INCOMPATIBILITIES

The simultaneous use of SACOX and certain antibiotic drugs (e.g. tiamulin) is contraindicated.

WITHDRAWAL PERIODS

Meat: zero (0) days

Eggs: DO NOT USE in birds laying eggs for human consumption or within 7 days of laying where eggs or egg products are to be used for human consumption or processing.

DISPOSAL

Shake and empty contents into medicated feed. Do not dispose of undiluted chemicals on site. Puncture or shred and bury containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt.

STORAGE CONDITIONS

Keep in dry place at temperature below 30°C.

SHELF LIFE AFTER INCORPORATION INTO COMPOUND FEEDINGSTUFF

3 months when stored at temperature up to 25°C (60% RH) or 1 (one) month when stored at temperature up to 40°C (75% RH).

PACKAGING AVAILABLE

Polyethylene-lined multiple-layer paper bags of 25 kg.

MANUFACTURER

BIOVET Joint Stock Company
 39 Petar Rakov Street, 4550 Peshtera,
 Bulgaria

MARKETING AUTHORIZATION HOLDER

Pahang Pharmacy Sdn. Bhd. (10826-K)
 Lot 5979 Jalan Teratai, 5½ Mile Off Jalan Meru,
 41050 Klang, Selangor Darul Ehsan, Malaysia

DATE OF REVISION

May 2018.