

25 mm



Rx Prescription only

Tadalafil STELLA

25 mm



- | | |
|-----|---|
| | <p>Name of the medicinal product</p> <p>Tadalafil STELLA 10 mg</p> <p>Tadalafil STELLA 20 mg</p> |
| 2. | <p>Special notice and recommendation</p> <p>Keep out of reach of children</p> <p>Read the package insert carefully before use</p> |
| 3. | <p>Qualitative and quantitative composition</p> <p>Each film-coated tablet Tadalafil STELLA 10 mg contains:</p> <p>Tadalafil.....10 mg</p> <p>Each film-coated tablet Tadalafil STELLA 20 mg contains:</p> <p>Tadalafil.....20 mg</p> |
| 4. | <p>Pharmaceutical form</p> <p>Film-coated tablet</p> <p>Tadalafil STELLA 10 mg:</p> <p>Yellow, oblong film - coated tablets, biconvex, engraved with logo "  " on one side and scored on the other side.</p> <p>Dimension: About 5.6 mm x 11.6 mm.</p> <p>Thickness: About 3.4 mm.</p> <p>The tablet can be divided into equal doses.</p> <p>Tadalafil STELLA 20 mg:</p> <p>Yellow, almond-shaped, film - coated tablet, biconvex, engraved with "T20" on one side and plain on the other side.</p> <p>Dimension: About 7.8 mm x 12.6 mm.</p> <p>Thickness: About 4.80 mm.</p> |
| 5. | <p>Indications</p> <p>Treatment of erectile dysfunction in adult males.</p> <p>In order for tadalafil to be effective, sexual stimulation is required.</p> <p>Tadalafil is not indicated for use by women.</p> |
| 6. | <p>Administration and dosage</p> <p>Adult men</p> <p>In general, the recommended dose is 10 mg taken prior to anticipated sexual activity and with or without food.</p> <p>In those patients in whom tadalafil 10 mg does not produce an adequate effect, 20 mg might be tried. It may be taken at least 30 minutes prior to sexual activity.</p> <p>The maximum dose frequency is once per day.</p> <p>Tadalafil 10 and 20 mg is intended for use prior to anticipated sexual activity and it is not recommended for continuous daily use.</p> <p>In patients who anticipate a frequent use of Tadalafil (i.e., at least twice weekly) a once daily regimen with the lowest doses of Tadalafil might be considered suitable, based on patient choice and the physician's judgement.</p> <p>In these patients the recommended dose is 5 mg taken once a day at approximately the same time of day. The dose may be decreased to 2.5 mg once a day based on individual tolerability.</p> <p>The appropriateness of continued use of the daily regimen should be reassessed periodically.</p> <p>Special populations</p> <p>Elderly men</p> <p>Dose adjustments are not required in elderly patients.</p> <p>Men with renal impairment</p> <p>Dose adjustments are not required in patients with mild to moderate renal impairment. For patients with severe renal impairment 10 mg is the maximum recommended dose. Once-a-day dosing of tadalafil is not recommended in patients with severe renal impairment.</p> <p>Men with hepatic impairment</p> <p>The recommended dose of Tadalafil is 10 mg taken prior to anticipated sexual activity and with or without food. There is limited clinical data on the safety of Tadalafil in patients with severe hepatic impairment (Child-Pugh Class C); if prescribed, a careful individual benefit/risk evaluation should be undertaken by the prescribing physician. There are no available data about the administration of doses higher than 10 mg of tadalafil to patients with hepatic impairment. Once-a-day dosing has not been evaluated in patients with hepatic impairment; therefore if prescribed, a careful individual benefit/risk evaluation should be undertaken by the prescribing physician.</p> <p>Men with diabetes</p> <p>Dose adjustments are not required in diabetic patients.</p> <p>Paediatric population</p> <p>There is no relevant use of Tadalafil in the paediatric population with regard to the treatment of erectile dysfunction.</p> |
| 7. | <p>Contraindications</p> <p>Hypersensitivity to the active substance or to any of the excipients in the drug products.</p> <p>Tadalafil was shown to augment the hypotensive effects of nitrates. This is thought to result from the combined effects of nitrates and tadalafil on the nitric oxide/cGMP pathway. Therefore, administration of tadalafil to patients who are using any form of organic nitrate is contraindicated.</p> <p>Tadalafil must not be used in men with cardiac disease for whom sexual activity is inadvisable. Physicians should consider the potential cardiac risk of sexual activity in patients with pre-existing cardiovascular disease.</p> <p>The use of tadalafil is contraindicated in the following groups of patients with cardiovascular disease:</p> <ul style="list-style-type: none"> + Patients with myocardial infarction within the last 90 days. + Patients with unstable angina or angina occurring during sexual intercourse. + Patients with New York Heart Association Class 2 or greater heart failure in the last 6 months. + Patients with uncontrolled arrhythmias, hypotension (< 90/50 mm Hg), or uncontrolled hypertension. + Patients with a stroke within the last 6 months. <p>Tadalafil is contraindicated in patients who have loss of vision in one eye because of non-arteritic anterior ischaemic optic neuropathy (NAION), regardless of whether this episode was in connection or not with previous PDE5 inhibitor exposure.</p> <p>The co-administration of PDE5 inhibitors, including tadalafil, with guanylate cyclase stimulators, such as riociguat, is contraindicated as it may potentially lead to symptomatic hypotension.</p> |
| 8. | <p>Special warnings and precautions for use</p> <p>Before treatment with tadalafil</p> <p>A medical history and physical examination should be undertaken to diagnose erectile dysfunction or benign prostatic hyperplasia and determine potential underlying causes, before pharmacological treatment is considered.</p> <p>Prior to initiating any treatment for erectile dysfunction, physicians should consider the cardiovascular status of their patients, since there is a degree of cardiac risk associated with sexual activity. Tadalafil has vasodilator properties, resulting in mild and transient decreases in blood pressure and as such potentiates the hypotensive effect of nitrates.</p> <p>The evaluation of erectile dysfunction should include a determination of potential underlying causes and the identification of appropriate treatment following an appropriate medical assessment. It is not known if tadalafil is effective in patients who have undergone pelvic surgery or radical non-nerve-sparing prostatectomy.</p> <p>Cardiovascular</p> <p>Serious cardiovascular events, including myocardial infarction, sudden cardiac death, unstable angina pectoris, ventricular arrhythmia, stroke, transient ischaemic attacks, chest pain, palpitations and tachycardia, have been reported either post marketing and/or in clinical trials. Most of the patients in whom these events have been reported had pre-existing cardiovascular risk factors. However, it is not possible to definitively determine whether these events are related directly to these risk factors, to tadalafil, to sexual activity, or to a combination of these or other factors.</p> <p>In patients who are taking alpha, blockers, concomitant administration of tadalafil may lead to symptomatic hypotension in some patients. The combination of tadalafil and doxazosin is not recommended.</p> <p>Vision</p> <p>Visual defects and cases of NAION have been reported in connection with the intake of tadalafil and other PDE5 inhibitors. Analyses of observational data suggest an increased risk of acute NAION in men with erectile dysfunction following exposure to tadalafil or other PDE5 inhibitors. As this may be relevant for all patients exposed to tadalafil, the patient should be advised that in case of sudden visual defect, he should stop taking tadalafil and consult a physician immediately.</p> <p>Decreased or sudden hearing loss</p> <p>Cases of sudden hearing loss have been reported after the use of tadalafil. Although other risk factors were present in some cases (such as age, diabetes, hypertension and previous hearing loss history) patients should be advised to stop taking tadalafil and seek prompt medical attention in the event of sudden decrease or loss of hearing.</p> <p>Hepatic impairment</p> <p>There is limited clinical data on the safety of single-dose administration of tadalafil in patients with severe hepatic insufficiency (Child-Pugh Class C). If tadalafil is prescribed, careful individual benefit/risk evaluation should be undertaken by the prescribing physician.</p> <p>Priapism and anatomical deformation of the penis</p> <p>Patients who experience erections lasting 4 hours or more should be instructed to seek immediate medical assistance. If priapism is not treated immediately, penile tissue damage and permanent loss of potency may result.</p> <p>Tadalafil should be used with caution in patients with anatomical deformation of the penis (such as angulation, cavernosal fibrosis, or Peyronie's disease) or in patients who have conditions which may predispose them to priapism (such as sickle cell anaemia, multiple myeloma or leukaemia).</p> <p>Use with CYP3A4 inhibitors</p> <p>Caution should be exercised when prescribing tadalafil to patients using potent CYP3A4 inhibitors (ritonavir, saquinavir, ketoconazole, itraconazole, and erythromycin), as increased tadalafil exposure (AUC) has been observed if the medicinal products are combined.</p> <p>Tadalafil and other treatments for erectile dysfunction</p> <p>The safety and efficacy of combinations of tadalafil and other PDE5 inhibitors or other treatments for erectile dysfunction have not been studied. The patients should be informed not to take tadalafil in such combinations.</p> <p>Lactose</p> <p>Tadalafil STELLA contains lactose. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take the medicinal product.</p> |
| 9. | <p>Pregnancy and lactation</p> <p>Tadalafil is not indicated for use by women.</p> <p>Pregnancy</p> <p>There are limited data from the use of tadalafil in pregnant women. Animal studies do not indicate direct or indirect harmful effects with respect to pregnancy, embryonal/foetal development, parturition or postnatal development. As a precautionary measure, it is preferable to avoid the use of tadalafil during pregnancy.</p> <p>Breastfeeding</p> <p>Available pharmacodynamic/toxicological data in animals have shown excretion of tadalafil in milk. A risk to the suckling child cannot be excluded. Tadalafil should not be used during breast feeding.</p> |
| 10. | <p>Effects on ability to drive and use machines</p> <p>Tadalafil has negligible influence on the ability to drive or use machines. Patients should be aware of how they react to tadalafil before driving or using machines.</p> |
| 11. | <p>Drug interactions</p> <p>Effects of other substances on tadalafil</p> <p>Cytochrome P450 inhibitors: Tadalafil is principally metabolised by CYP3A4. A selective inhibitor of CYP3A4, ketoconazole, increased tadalafil exposure (AUC) and C_{max} relative to the AUC and C_{max} values for tadalafil alone. Ritonavir, a protease inhibitor (200 mg twice daily), which is an inhibitor of CYP3A4, CYP2C9, CYP2C19, and CYP2D6, increased tadalafil exposure (AUC) with no change in C_{max}. Although specific interactions have not been studied, other protease inhibitors, such as saquinavir, and other CYP3A4 inhibitors, such as erythromycin, clarithromycin, itraconazole, and grapefruit juice, should be coadministered with caution, as they would be expected to increase plasma concentrations of tadalafil. Consequently, the incidence of the adverse reactions might be increased.</p> <p>Transporters: The role of transporters (for example, p-glycoprotein) in the disposition of tadalafil is not known. Therefore, there is the potential of drug interactions mediated by inhibition of transporters.</p> <p>Cytochrome P450 inducers: A CYP3A4 inducer, rifampicin, reduced tadalafil AUC relative to the AUC values for tadalafil alone (10 mg). This reduced exposure can be anticipated to decrease the efficacy of tadalafil; the magnitude of decreased efficacy is unknown. Other inducers of CYP3A4, such as phenobarbital, phenytoin, and carbamazepine, may also decrease plasma concentrations of tadalafil.</p> |

