

## TAIKOSOL 300MG/ML SOLUTION

### Description and Composition

TAIKOSOL 300MG/ML SOLUTION is a light yellow to yellow clear solution. Each ml contains 300mg of Florfenicol as its active ingredient.

### Pharmacodynamics

Like chloramphenicol, florfenicol is a broad-spectrum antibiotic that has activity against many bacteria. It acts by binding to the 50S ribosome, thereby inhibiting bacterial protein synthesis.

### Pharmacokinetics

During a continuous treatment of broiler chicken with florfenicol in drinking water at a daily dose of 26mg/kg bw for 18 hours for 3 days, the estimated mean serum concentration of florfenicol was 711µg/l. No florfenicol was detected in the serum, 72 hours after the terminal dose.

In pigs, during a continuous treatment by florfenicol via drinking water at daily dose of approximately 16mg/kg bw for 5 days, the serum concentrations of florfenicol ranged from 1500 to 2500µg/l.

### Indication

Chickens: Prevention and treatment of bacterial infections associated with *Salmonella* spp. and *E. coli* susceptible to florfenicol.

Pigs: Prevention and treatment of bacterial infections associated with *Actinobacillus pleuropneumoniae*, *Pasteurella multocida*, *Mycoplasma pneumoniae* (Mycoplasma) and other respiratory diseases, *Salmonella typhimurium*, and *Streptococcus suis* susceptible to florfenicol.

### Recommended Dose

Chickens: 0.067ml (equivalent to 20mg Florfenicol) per kg body weight for 5 days. This dosage rate is equivalent to 33.34ml per 100L of drinking water for 5 days.

Pigs: 0.34ml (equivalent to 100mg Florfenicol) per 10kg body weight for 5 days. This dosage rate is equivalent to 33.34ml per 100L of drinking water for 5 days.

### Route of Administration

For oral administration via drinking water.

### Contraindication

None known.

### Warning and Precautions

Special warning for target species:

Do not use in animals that have experienced shock or sensitivity to product.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

- 1) Do not use chlorinated water during dosing.
- 2) Do not use a galvanized bucket.
- 3) Can cause precipitation when diluted at a ratio other than as prescribed (high concentration).
- 4) Mixture of product with water should be mixed homogeneously.

Other Precautions:

- 1) This product is a veterinary medicinal product, do not use it in humans.
- 2) Use according to the instructions of the veterinarian.
- 3) Read this instruction manual carefully before use.
- 4) Safety and efficacy have not been established except for the indicated species, therefore, do not use it arbitrarily.
- 5) Persons handling the product should wear gloves, masks, protective clothing to prevent direct skin

contact. Avoid breathing in.

- 6) In case of contact with skin or eyes when handling chemicals, immediately wash off and observe abnormal symptoms. Seek medical attention immediately.

#### **Interaction with Other Medicaments**

No specific drug interactions for florfenicol were located, but the drug may behave similarly to chloramphenicol. If so, florfenicol could antagonize the bactericidal activity of the penicillins or aminoglycosides. This antagonism has not been demonstrated *in vivo*, and these drug combinations have been used successfully many times clinically. Other antibiotics that bind to the 50S ribosomal subunit of susceptible bacteria (erythromycin, clindamycin, lincomycin, tylosin, etc.) may potentially antagonize the activity of chloramphenicol or vice versa, but the clinical significance of this potential interaction has not been determined. For other drug interactions that florfenicol may share with chloramphenicol, see the monograph for chloramphenicol or refer to other drug information resources.

#### **Pregnancy and Lactation**

Do not use in laying hens.

#### **Side Effects**

In continuous dosing, temporary inflammation may occur in the cloaca of the chicken and in the anus of the pig.

#### **Symptoms and Treatment of Overdose**

In toxicology studies where feeder calves were injected with up to 10 times the recommended dosage, the adverse effects noted above were seen, plus increased serum enzymes. These effects were generally transient in nature. Long-term (43 days) standard dosage studies showed a transient decrease in feed consumption, but no long-term negative effects were noted.

#### **Storage Condition**

Store below 30°C.

#### **Shelf life**

3 years.

After first opening the container, any product remaining after 24 hours should be discarded.

To be used within 24 hours after dilution.

#### **Withdrawal Periods**

Chickens: 5 days

Pigs: 16 days

#### **Packing**

200ml, 500ml and 1L

Revised date: 3.12.18

|                                                                                     |                                                                                                                                                                                              |
|-------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|  | Manufacturer & Product Registration Holder                                                                                                                                                   |
|                                                                                     | Range Pharma Sdn Bhd<br>No. 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia<br>Tel: 603-61568708 Fax: 603-61568707<br>Email: info@rangepharma.com |