

GDSGMY BuzzAway Bite Relief Cream_Leaflet

Size: 100 (W) x 160 (H) mm

GD875LR2M

Front Panel

guardian

BUZZAWAY BITE RELIEF CREAM

Active ingredients: Camphor 5.83%w/w, Menthol 3.92%w/w, Methyl Salicylate 3.75%w/w.
Preservative: Chloracresol 0.2%w/w.
Product description: White colour, smooth application, sticky and characteristic odor cream.
Pharmacodynamics:
Camphor
Camphor is a stimulant, rubefacient, antipruritic and feebly antiseptic. When in combination with other ingredient like menthol, methyl salicylate, it becomes ideal for neurogia and other painful areas.
Menthol
Menthol dilates the blood vessels causing a sensation of coldness, followed by an analgesic effect. Menthol also acts as a penetration enhancer, increasing the penetrational drugs when applied on the skin, to give a faster onset of action.
Methyl Salicylate
Methyl salicylate is a salicylic acid derivative. Salicylates inhibit cyclooxygenase, thereby reducing the formation of prostaglandins and cause platelet dysfunction. Methyl salicylate is used topically as a counter irritant. Upon application, it is absorbed through the skin and is applied for the relief of rheumatic pain and muscle or joints pain.

Pharmacokinetics:
Camphor
Camphor is readily absorbed from all administration sites. Camphor is well absorbed after dermal exposure. It is hydroxylated in the liver to yield hydroxycamphor metabolites which are then conjugated with glucuronic acid and excreted in the urine. Camphor crosses the placenta.
Menthol
After absorption, menthol is excreted in the urine and bile as a glucuronide.
Methyl Salicylate
May be absorbed through intact skin. The absorption of topical salicylates is proportional to the surface area involved, duration of exposure, concentration and skin integrity. Absorption characteristics of salicylates vary with the dose, formulation and route of administration. Percutaneous absorption is enhanced by exercise, heat, occlusion or disruption of the integrity of the skin or application to large areas of skin. Both the rate and extent of absorption increases after repeated application, increasing the bioavailability. Methyl salicylate is extensively metabolised to salicylic acid in the dermal and subcutaneous tissues after topical application. At therapeutic levels, the half life of salicylates is 2 - 4 hours. As salicylate level increase into the toxic range, the half-life can be greater than 18 hours.

Indication: Soothes and relieves itch and swelling from mosquito and insect bites.
Direction for use: Apply gently onto affected area. Re-apply as necessary. Wash hands thoroughly before and after each application.
Route of administration: Topical.

Contraindications: Patient with known hypersensitivity to any of the ingredients in the product.
Warnings and precautions: Can cause convulsions. This product is contraindicated in children below 2 years of age. Caution must be exercised when older children are treated. It is dangerous to place any camphor containing product into the mouth of children. A small amount applied this way may cause immediate collapse. This product contains methyl salicylate and when applied or rub

Manufactured under the commission of DB Brands Limited
5/F Deep House, Takko Place, Quarry Bay, Hong Kong
Distributed By:
Guardian Health & Beauty :
21 Tampines North Drive 2, #03-01, Giant,
Singapore 528765
Tel: (65) 6694 6400
Malaysia Product Registration Holder
Guardian Health & Beauty Sdn. Bhd. (201407024953)
Level 11 & First Avenue, Bandar Utama,
47800, Petaling Jaya,
Selangor Darul Ehsan, Malaysia.
Tel: (603) 3 2777 7700
Guardian Health & Beauty Sdn. Bhd. (RC00002394)
Lot 58865 Rimba Point, Kampong Rimba,
Mukim Gadong BE3119, Brunei Darussalam

Manufactured By:
Sunkist Wen Chai Ong Sdn Bhd
24, Jalan Lambak, Taman Johor,
81200 Johor Bahru, Johor, Malaysia.
MA11903015XC
Date of revision: Dec 2024
GD875LR2M

guardian

蚊虫叮咬止痒膏

活性成分: 薄荷 5.83%w/w, 薄荷醇 3.92%w/w, 水杨酸甲酯 3.75%w/w.
辅料: 硬脂醇 0.2 %w/w, 硬脂醇 0.2 %w/w.
产品描述: 白色, 涂感平滑, 带有特奇气味的粘稠膏膏。
适应症:
本品是一种兴奋剂, 发汗剂, 止痛剂和生物防腐剂。当与薄荷醇, 水杨酸甲酯等其他成分配合使用时, 它成为神经痛和其他疼痛部位的理想选择。
薄荷醇
薄荷醇能使血管扩张并引起寒冷却, 并起镇痛作用。薄荷醇还可以作为渗透促进剂, 增加药物在皮肤上的穿透力, 从而加快了药物的起效时间。

on the skin, can be absorbed through the skin into blood. For patients taking warfarin, excessive application on to the skin for muscle or joint pains may increase the chances of bleeding. Avoid contact With the eyes and mucous membrane. Do not apply to open wounds or damaged skin. Avoid excessive use during pregnancy and lactation. Do not apply over large areas of the body or bandage or use with heating pads. Initially apply a small amount to a limited skin area and observe whether there is discomfort or itch before applying to a larger skin area. Discontinue use if irritation develops. For itchy skin, only use if irritation occurs and seek medical advice immediately. Keep out of reach of children.
Effects on ability to drive and use machines: Not known.

Interactions with other medications: For patients taking warfarin, excessive application on to the skin for muscle or joints pains may increase the chances of bleeding.
Statement on usage during pregnancy and lactation: Safety for use in pregnancy has not been established therefore the potential benefit of the product should be weighed against the possible risks to the mother and child.

Adverse effects/undesirable effects: Headache, dizziness, nausea and vomiting, skin irritation, contact dermatitis, rash, itching, redness or swelling, burning or stinging sensation may occur. May cause hypersensitivity allergic reactions in some individuals with sensitive skin.
Symptoms and treatment of overdose:
Symptoms of methyl salicylate overdose
Salicylate intoxication can occur after ingestion or topical application of methyl salicylate. Mild chronic salicylate intoxication or salicylism, usually occurs only after repeated use of large doses. Salicylism can also occur following excessive topical application of salicylates. Symptoms include dizziness, tinnitus, deafness, sweating, nausea and vomiting, headache and confusion and may be controlled by reducing the dosage. Symptoms of more severe intoxication or of acute poisoning following overdose include hyperventilation, fever, restlessness, ketosis and respiratory alkalosis and metabolic acidosis. Depression of the CNS may lead to coma; cardiovascular collapse and respiratory failure may also occur.

Symptoms of camphor/ menthol overdose
Ingestion of significant quantities is reported to cause symptom: severe abdominal pain, nausea, vomiting, vertigo, ataxia, drowsiness and coma; instant collapse in infants after the local application of menthol or camphor to their nostrils. Seizures may be the first clinical sign of severe toxicity of camphor; however, seizures are usually self- limited. Severe toxicity of camphor can result in delirium, visual hallucinations, cerebral edema and status epilepticus. Systemic toxicity may include hypotension, tachycardia, respiratory failure and death.
Treatment
The stomach should be emptied by gastric lavage or administration of oral activated charcoal. Fluid and electrolyte management is the mainstay of treatment with the immediate aim of correction of acidosis, hyperpyrexia, hypokalaemia and dehydration if present.

Storage conditions: Store below 30°C, away from direct sunlight.
Shelf life: Please refer to the packaging. Do not use beyond the expiry date.
Dosage form: Cream. **Packing:** 30g in aluminum tube.
The information contained in this leaflet is limited. For further information, please consult your doctor or pharmacist.

Manufactured By:
Sunkist Wen Chai Ong Sdn Bhd
24, Jalan Lambak, Taman Johor,
81200 Johor Bahru, Johor, Malaysia.
MA11903015XC
Date of revision: Dec 2024
GD875LR2M

Manufactured By:
Sunkist Wen Chai Ong Sdn Bhd
24, Jalan Lambak, Taman Johor,
81200 Johor Bahru, Johor, Malaysia.
MA11903015XC
Date of revision: Dec 2024
GD875LR2M

水杨酸甲酯
水杨酸甲酯是一种水杨酸衍生物。水杨酸酯能抑制环氧合酶, 从而减少前列腺素的形成而引起血小板功能障碍。局部使用水杨酸酯可能为反射刺激。使用后, 它通过皮肤进入血液循环并迅速在肌肉和组织内下降。
现代动力学:
薄荷醇很容易从所有途径吸收。水杨酸酯到皮肤后很容易被吸收。它在肝脏中被羟基化, 产生羟基水杨酸代谢物, 然后与葡萄糖醛酸结合并通过尿液排出。薄荷醇可穿过胎盘。
薄荷醇
薄荷醇能使血管扩张并引起寒冷却, 并起镇痛作用。薄荷醇还可以作为渗透促进剂, 增加药物在皮肤上的穿透力, 从而加快了药物的起效时间。

水杨酸甲酯
水杨酸甲酯是一种水杨酸衍生物。水杨酸酯能抑制环氧合酶, 从而减少前列腺素的形成而引起血小板功能障碍。局部使用水杨酸酯可能为反射刺激。使用后, 它通过皮肤进入血液循环并迅速在肌肉和组织内下降。
现代动力学:
薄荷醇很容易从所有途径吸收。水杨酸酯到皮肤后很容易被吸收。它在肝脏中被羟基化, 产生羟基水杨酸代谢物, 然后与葡萄糖醛酸结合并通过尿液排出。薄荷醇可穿过胎盘。
薄荷醇
薄荷醇能使血管扩张并引起寒冷却, 并起镇痛作用。薄荷醇还可以作为渗透促进剂, 增加药物在皮肤上的穿透力, 从而加快了药物的起效时间。

水杨酸甲酯
水杨酸甲酯是一种水杨酸衍生物。水杨酸酯能抑制环氧合酶, 从而减少前列腺素的形成而引起血小板功能障碍。局部使用水杨酸酯可能为反射刺激。使用后, 它通过皮肤进入血液循环并迅速在肌肉和组织内下降。
现代动力学:
薄荷醇很容易从所有途径吸收。水杨酸酯到皮肤后很容易被吸收。它在肝脏中被羟基化, 产生羟基水杨酸代谢物, 然后与葡萄糖醛酸结合并通过尿液排出。薄荷醇可穿过胎盘。
薄荷醇
薄荷醇能使血管扩张并引起寒冷却, 并起镇痛作用。薄荷醇还可以作为渗透促进剂, 增加药物在皮肤上的穿透力, 从而加快了药物的起效时间。

水杨酸甲酯
水杨酸甲酯是一种水杨酸衍生物。水杨酸酯能抑制环氧合酶, 从而减少前列腺素的形成而引起血小板功能障碍。局部使用水杨酸酯可能为反射刺激。使用后, 它通过皮肤进入血液循环并迅速在肌肉和组织内下降。
现代动力学:
薄荷醇很容易从所有途径吸收。水杨酸酯到皮肤后很容易被吸收。它在肝脏中被羟基化, 产生羟基水杨酸代谢物, 然后与葡萄糖醛酸结合并通过尿液排出。薄荷醇可穿过胎盘。
薄荷醇
薄荷醇能使血管扩张并引起寒冷却, 并起镇痛作用。薄荷醇还可以作为渗透促进剂, 增加药物在皮肤上的穿透力, 从而加快了药物的起效时间。

水杨酸甲酯
水杨酸甲酯是一种水杨酸衍生物。水杨酸酯能抑制环氧合酶, 从而减少前列腺素的形成而引起血小板功能障碍。局部使用水杨酸酯可能为反射刺激。使用后, 它通过皮肤进入血液循环并迅速在肌肉和组织内下降。
现代动力学:
薄荷醇很容易从所有途径吸收。水杨酸酯到皮肤后很容易被吸收。它在肝脏中被羟基化, 产生羟基水杨酸代谢物, 然后与葡萄糖醛酸结合并通过尿液排出。薄荷醇可穿过胎盘。
薄荷醇
薄荷醇能使血管扩张并引起寒冷却, 并起镇痛作用。薄荷醇还可以作为渗透促进剂, 增加药物在皮肤上的穿透力, 从而加快了药物的起效时间。

guardian

KRIM KELEGAAN GITAN

Komposisi: Camphor 5.83%w/w, Menthol 3.92%w/w, Methyl Salicylate 3.75%w/w.
Bahan Pengawet: Chloracresol 0.2%w/w.

Deskripsi Produk: Krem berwarna putih, mudah disapu, melekat dan mempunyai bau citran.

Farmakodinamik:
Camphor
Camphor adalah perangsang, rubefasiens, antipruritik dan antiseptik lemah. Apabila digabungkan dengan ramuan lain seperti menthol, methyl salicylate, ia menjadi sesuai untuk neurogia dan kawasan-kawasan lain yang sakit.
Menthol
Menthol dilat saliran darah dan menyebabkan sensasi kesejukan, diikuti dengan kesan analgesik. Menthol juga bertindak sebagai penembusan, meningkatkan penembusan ubat-ubatan apabila digunakan pada kulit, untuk memberi permulaan tindakan yang lebih cepat.

Methyl Salicylate
Methyl Salicylate adalah derivatif asid salisilik. Salicylate menghambat siklooksigenase, dengan itu mengurangkan pembentukan prostaglandin dan menyebabkan distensi platelet. Methyl Salicylate digunakan secara topikal sebagai perangsang kawan. Selepas aplikasi, ia diserap melalui kulit dan digunakan untuk melegakan kesakitan reumatik dan kesakitan otot atau sendi.
Farmakokinetik:
Camphor
Camphor mudah diserap dari semua laluan pemberian. Camphor diserap dengan baik selepas pendedahan derma. Ia dihidrolisis di hati untuk menghasilkan metabolit hydroxycamphor yang kemudiannya berkonjugat dengan asid glucuronic dan diekskresikan dalam urin. Camphor melepasi plasenta.
Menthol
Selepas penyerapan, menthol diekskresikan dalam urin dan hempedu sebagai glukuronid.
Methyl Salicylate
Boleh diserap melalui kulit uluh. Penyerapan salicylate topikal adalah berkadar dengan kawasan permukaan yang terbitar, tempoh dedahan, kepekatan dan integriti kulit. Ciri-ciri penyerapan salicylate berbeza dengan dos, permukaan laluan pemberian uluh. Penyerapan perkutanus diperingkatkan dengan senaman, haba, aktiviti atau gangguan keuluhan kulit atau apabila ke kawasan kulit yang besar. Dua-dua-dua kadar dan tahap penyerapan meningkat selepas aplikasi berulang, meningkatkan biokefektifan. Methyl Salicylate dimetabolisme secara meluas kepada asid salisilik dalam isu kulit dan subkutin selepas pemohonan topikal. Pada tahap terapeutik, separuh hayat salicylate adalah 2 - 4 jam. Apabila paras salicylate meningkat ke dalam julat toksik, separuh hayat boleh lebih besar daripada 18 jam.

Indikasi: Untuk menenang dan melegakan kegelatan dan bengkak yang disebabkan oleh gigitan nyamuk dan serangga.
Dos saranan: Sapukan krim pada bahagian yang berkenaan dengan lembut. Dilutangi sekiranya perlu. Basuhkan tangan secara menyeluruh sebelum dan selepas guna.
Laluan pemberian ubat: Topikal.

Kontraindikasi: Pesakit yang diketahui hipersensitif kepada mana-mana ramuan dalam produk.
Amaran dan perhatian: Boleh menyebabkan konvulsi. Ubat ini tidak sesuai untuk bayi di bawah umur 2 tahun dan perhatian perlu diberikan kepada kanak-kanak berumur 2 tahun ke atas apabila menggunakannya. Sapu ubat yang mengandungi camphor pada hidung kanak-kanak adalah berbahaya kerana boleh menyebabkan pengsan. Ubat ini mengandungi methyl salicylate dan ia mungkin boleh diserap masuk ke dalam darah melalui kulit. Penayapan yang berlebihan oleh pesakit yang mengambil ubat ini boleh meningkatkan risiko pendarahan. Tidak dibenarkan.
Interaksi dengan ubat lain: Untuk pesakit yang mengambil warfarin, aplikasi berlebihan pada kulit untuk kesakitan otot atau sendi akan meningkatkan peluang pendarahan.
Kehamilan dan penyusuan: Sila rujuk kepada penderangan Bahasa Inggeris. Keselamatan untuk kegunaan semasa kehamilan belum dimantapkan. Oleh itu potensi manfaat produk terhadap kemungkinan risiko kepada ibu dan anak harus ditimbang.
Kesan sampingan: Kulit kepala, pening, loya dan muntah, kelegaan kulit, dermatitis sentuh, ruam, kegelatan, kemerahan atau pembengkakan, pembakaran atau rasa menyengat mungkin berlaku. Boleh menyebabkan tindak balas hipersensitifasi/dergi pada sesetengah individu yang mempunyai masalah kulit sensitif.

Simpulan dan rawatan untuk kelebihan dos:
Simpulan dos berlebihan methyl salicylate
Intoksikasi salicylate boleh berlaku selepas pengesanan atau aplikasi topikal methyl salicylate. Keracunan salicylate kronik yang ringan atau salisimisme biasanya berlaku hanya selepas penggunaan berulang dalam dos yang besar. Salisimisme juga boleh berlaku akibat aplikasi topikal salicylate yang berlebihan. Simptom termasuk kepening, inisius, kepekatan, berpeluh, loya dan muntah, sakit kepala dan kekeliruan dan boleh dikawal dengan mengurangkan dos. Simptom keracunan yang lebih teruk atau keracunan akut diikuti dengan dos berlebihan termasuk hiperventilasi, demam, keresahan, ketosis dan alkalosis respiratori dan asidosis metabolik. Depresi SSF boleh menyebabkan koma. Kematian kardiovaskular dan kegagalan pernafasan juga mungkin berlaku.
Simpulan dos berlebihan camphor/ menthol
Pengambilan kuantiti yang banyak telah dilaporkan boleh menyebabkan simptom seperti sakit perut yang teruk, loya, muntah, vertigo, ataksia, mengikut "dan koma". Bayi akan pengsan segera selepas aplikasi menthol atau camphor ke lubang hidung mereka.
Penyitaan mungkin adalah tanda klinikal yang pertama untuk ketoksikan camphor yang teruk. Walau bagaimanapun, penyitaan biasanya terhad kepada diri sendiri. Ketoksikan yang teruk boleh mengakibatkan delirium, halusinasi visual, edema serebrum dan status epilepticus. Ketoksikan sistemik termasuk hipotensi, takikardia, kegagalan pernafasan dan kematian.
Rawatan
Perlu harus dikawalskan dengan loya gaster atau pengambilan arang diaktifkan secara oral. Pengurusan bendalir dan elektrolit adalah sangat rawatan untuk pembetulan asidosis, hiperelektrolit, hipokalemia dan penyahhidrat jika hadir.
Kedudukan penyimpanan: Simpan di bawah 30°C, jauhkan daripada pancaran terus matahari.
Tempoh hayat: Sila rujuk kepada bungkusan. Jangan guna selepas tarikh luput.
Dos: Krim. **Sila bungkusan:** 30g dalam tiub aluminium.
Maklumat dalam risalah ini adalah terhad. Untuk informasi selanjutnya, sila jumpa doktor atau ahli farmasi.

Kelebihan dos berlebihan:
Methyl salicylate overdose
Salicylate intoxication can occur after ingestion or topical application of methyl salicylate. Mild chronic salicylate intoxication or salicylism, usually occurs only after repeated use of large doses. Salicylism can also occur following excessive topical application of salicylates. Symptoms include dizziness, tinnitus, deafness, sweating, nausea and vomiting, headache and confusion and may be controlled by reducing the dosage. Symptoms of more severe intoxication or of acute poisoning following overdose include hyperventilation, fever, restlessness, ketosis and respiratory alkalosis and metabolic acidosis. Depression of the CNS may lead to coma; cardiovascular collapse and respiratory failure may also occur.
Symptoms and treatment of overdose:
Symptoms of methyl salicylate overdose
Salicylate intoxication can occur after ingestion or topical application of methyl salicylate. Mild chronic salicylate intoxication or salicylism, usually occurs only after repeated use of large doses. Salicylism can also occur following excessive topical application of salicylates. Symptoms include dizziness, tinnitus, deafness, sweating, nausea and vomiting, headache and confusion and may be controlled by reducing the dosage. Symptoms of more severe intoxication or of acute poisoning following overdose include hyperventilation, fever, restlessness, ketosis and respiratory alkalosis and metabolic acidosis. Depression of the CNS may lead to coma; cardiovascular collapse and respiratory failure may also occur.

Symptoms of camphor/ menthol overdose
Ingestion of significant quantities is reported to cause symptom: severe abdominal pain, nausea, vomiting, vertigo, ataxia, drowsiness and coma; instant collapse in infants after the local application of menthol or camphor to their nostrils. Seizures may be the first clinical sign of severe toxicity of camphor; however, seizures are usually self- limited. Severe toxicity of camphor can result in delirium, visual hallucinations, cerebral edema and status epilepticus. Systemic toxicity may include hypotension, tachycardia, respiratory failure and death.
Treatment
The stomach should be emptied by gastric lavage or administration of oral activated charcoal. Fluid and electrolyte management is the mainstay of treatment with the immediate aim of correction of acidosis, hyperpyrexia, hypokalaemia and dehydration if present.
Storage conditions: Store below 30°C, away from direct sunlight.
Shelf life: Please refer to the packaging. Do not use beyond the expiry date.
Dosage form: Cream. **Packing:** 30g in aluminum tube.
The information contained in this leaflet is limited. For further information, please consult your doctor or pharmacist.

Manufactured By:
Sunkist Wen Chai Ong Sdn Bhd
24, Jalan Lambak, Taman Johor,
81200 Johor Bahru, Johor, Malaysia.
MA11903015XC
Date of revision: Dec 2024
GD875LR2M

Manufactured By:
Sunkist Wen Chai Ong Sdn Bhd
24, Jalan Lambak, Taman Johor,
81200 Johor Bahru, Johor, Malaysia.
MA11903015XC
Date of revision: Dec 2024
GD875LR2M

Manufactured By:
Sunkist Wen Chai Ong Sdn Bhd
24, Jalan Lambak, Taman Johor,
81200 Johor Bahru, Johor, Malaysia.
MA11903015XC
Date of revision: Dec 2024
GD875LR2M

Manufactured By:
Sunkist Wen Chai Ong Sdn Bhd
24, Jalan Lambak, Taman Johor,
81200 Johor Bahru, Johor, Malaysia.
MA11903015XC
Date of revision: Dec 2024
GD875LR2M

Manufactured By:
Sunkist Wen Chai Ong Sdn Bhd
24, Jalan Lambak, Taman Johor,
81200 Johor Bahru, Johor, Malaysia.
MA11903015XC
Date of revision: Dec 2024
GD875LR2M

Manufactured By:
Sunkist Wen Chai Ong Sdn Bhd
24, Jalan Lambak, Taman Johor,
81200 Johor Bahru, Johor, Malaysia.
MA11903015XC
Date of revision: Dec 2024
GD875LR2M

Manufactured By:
Sunkist Wen Chai Ong Sdn Bhd
24, Jalan Lambak, Taman Johor,
81200 Johor Bahru, Johor, Malaysia.
MA11903015XC
Date of revision: Dec 2024
GD875LR2M