

HemoRid CT

MICRONISED TABLET

COMPOSITION

Each film-coated tablet contains Micronised Diosmin 450 mg and Micronised Hesperidin 50 mg.

DESCRIPTION

Oval shape, film-coated pinkish brown tablet.

PHARMACODYNAMICS

Hemo Rid CT contains the Bioflavonoids, Diosmin and Hesperidin, which are thought to improve capillary function by reducing abnormal leakage. Hemo Rid CT reinforces venous tone by prolonging the activity of parietal noradrenaline, thereby decreasing venous capacitance, venous distensibility and venous emptying time. It also inhibits the release of mediators e.g. free radicals and prostaglandins, thus normalising capillary permeability and strengthening capillary resistance. Hemo Rid CT also improves lymphatic drainage by increasing lump flow and lymph oncotic pressure, producing phlebotonic and vasculo protective effects which significantly improve the chronic symptoms of hemorrhoidal disease.

PHARMACOKINETICS

In humans, following oral administration of Diosmin:

- Excretion is essentially faecal and urinary excretion is on average 14% of the administered quantity.
- The elimination half life is of 11 hours.
- The product is highly metabolised, this metabolism is revealed by the presence of different phenol acids in the urine.

INDICATIONS

For the treatment of functional symptoms related to acute hemorrhoidal attacks. Treatment of symptoms related to venolymphatic insufficiency (heavy legs, pain, early morning restless legs).

DOSAGE AND ADMINISTRATION

The dosage of Hemo Rid CT will depend on the severity of the disease.

Chronic Haemorrhoids: 2 tablets daily in two divided doses, midday and evening at meal times.

Acute Hemorrhoidal Attack: 6 tablets daily in 2 divided doses, for 4 days and then 4 tablets daily in 2 divided doses for 3 days.

(Tablets are to be taken after meals)

Chronic Venous Insufficiency: 2 tablets daily in two divided doses, midday and evening at meal times.

Route of Administration

Oral

CONTRAINDICATION

Patients with known hypersensitivity to Diosmin or Hesperidin, or to any of the components of the product.

WARNINGS AND PRECAUTIONS FOR USE

Hemorrhoidal attacks:

The administration of this product does not preclude treatment for other anal conditions.

The treatment must be short term. If symptoms do not subside promptly, a proctological examination should be performed and the treatment should be reviewed.

Use in pregnancy

Studies in animals are inadequate or may be lacking, but available data from animal studies show no evidence of teratogenic effect. Moreover, no harmful effects have been reported to date in humans.

Breast-feeding

In the absence of data on excretion into breast milk, breastfeeding is not recommended during treatment.

INTERACTIONS WITH OTHER MEDICAMENTS

No interaction studies have been performed. However, taking into consideration the huge post marketing experience of Diosmin and Hesperidin, no drug interaction has been reported to date.

UNDESIRABLE EFFECTS

Some cases of minor gastrointestinal and autonomic disorders have been reported (diarrhoea, dyspepsia, nausea, vomiting), which never required cessation of treatment.

Tabulated list of Adverse Reactions

System Organ Class	Frequency	Preferred Term
Nervous system disorders	Rare	Dizziness
		Headache
		Malaise
Gastrointestinal disorders	Common	Diarrhoea
		Dyspepsia
		Nausea
		Vomiting
	Uncommon	Colitis
Skin and subcutaneous tissue disorders	Rare	Pruritus
		Rash
		Urticaria

210mm



100% Black

148mm

210mm

OVERDOSAGE

There are no known cases of overdose with Hemo Rid CT. There is no specific antidote for Diosmin and Hesperidin. Treatment should be symptomatic should excessive quantities be ingested.

PRESENTATION

1 x 10's film-coated tablets in blister packs
3 x 10's film-coated tablets in blister packs
10 x 10's film-coated tablets in blister packs

STORAGE CONDITIONS

- STORE IN PLACE BELOW 30°C.
- PROTECT FROM LIGHT.
- KEEP OUT OF REACH OF CHILDREN.

SHELF LIFE

Please refer to outer package

Product Registration Holder :

**DUOPHARMA MANUFACTURING
(BANGI) SDN. BHD.**

Lot No. 2, 4, 6, 8 & 10, Jalan P/7, Section 13,
Bangi Industrial Estate, 43650 Bandar Baru Bangi,
Selangor, Malaysia.

Manufacturer:

**DUOPHARMA MANUFACTURING
(BANGI) SDN. BHD.**

Lot No. 2 & 4, Jalan P/7, Section 13,
Bangi Industrial Estate, 43650 Bandar Baru Bangi,
Selangor, Malaysia.

Revision Date: Apr 2025
150000XXXX N.X



100% Black