

BOFLOX 100 mg/ml

Solution for injection Marbofloxacin

Composition

1 ml contains: **Active ingredient:** Marbofloxacin, 100 mg.
Excipients: Metacresol, 2 mg; Excipient, q.s.

Yellow greenish to yellow brownish, clear solution.

Indications

Cattle: Treatment of respiratory infections caused by sensitive strains of *Pasteurella multocida*, *Mannheimia (Pasteurella) haemolytica* and *Mycoplasma bovis*. Treatment of acute mastitis caused by *E. coli* strains sensitive to marbofloxacin during the lactation period.

Pigs (sows): Treatment of Metritis Mastitis Agalactia Syndrome caused by bacterial strains sensitive to marbofloxacin.

Contraindications

Do not use in bacterial infections with resistance to other fluoroquinolones (cross resistance). Do not use in animals previously found to be hypersensitive to marbofloxacin or other quinolone.

Adverse reactions

Administration by intramuscular route may cause transient local reactions such as pain and swelling at the injection site and inflammatory lesions which persist, at least, 12 days after injection. However, in cattle, subcutaneous route was shown to be better tolerated locally than by intramuscular route. Therefore, the subcutaneous route is recommended in heavy cattle. For the injections, the neck should be preferred in cattle and pigs. No other undesirable effects have been observed in cattle and pigs.

Target species

Cattle and pigs (sows).

Dosage for each species, routes and method of administration

Cattle: Intramuscular, subcutaneous or intravenous use. Treatment of respiratory infections caused by sensitive strains of *Pasteurella multocida*, *Mannheimia haemolytica*: The recommended dose is 8 mg marbofloxacin/kg bw (2 ml product/25 kg bw) in a single intramuscular injection. Treatment of respiratory infections caused by sensitive strains of *Mycoplasma bovis*: The recommended dose is 2 mg marbofloxacin/kg bw (1 ml product/50 kg bw) in a single daily injection by SC or IM route, for 3 to 5 days. The first injection may also be given by the IV route. Acute mastitis: The recommended dose is 2 mg marbofloxacin/kg bw (1 ml product/50 kg bw) in a single daily

injection by subcutaneous or intramuscular route, for 3 consecutive days. The first injection may also be given by the IV route.

Pigs (sows): Intramuscular use.

The recommended dose is 2 mg marbofloxacin/kg bw (1 ml product/50 kg bw) in a single daily injection, for 3 consecutive days.

Withdrawal period

Cattle:

Indication: Respiratory infections.

Dose: 2 mg/kg for 3 to 5 days (IV/IM/SC).

Meat: 6 days.

Milk: 36 hours.

Dose: 8 mg/kg on a single occasion (IM).

Meat: 3 days.

Milk: 72 hours.

Indication: Mastitis.

Dose: 2 mg/kg for 3 days (IV/IM/SC).

Meat: 6 days.

Milk: 36 hours.

Pigs (sows): Meat: 4 days.

Storage precautions

Keep out of the sight and reach of children.

Do not store above 30 °C. Protect from light.

Once opened, use within 28 days stored below 25 °C and 60 ± 5 % RH.

Special warnings

Special warnings for each target species: None.

Special precautions for use in animals: Fluoroquinolones should be reserved for the treatment of clinical conditions which have responded poorly, or are expected to respond poorly, to other classes of antimicrobials. Whenever possible, fluoroquinolones should only be used based on susceptibility testing. Official and local antimicrobial policies should be taken into account when the product is used. Use of the product deviating from the instructions may increase the prevalence of bacteria resistant to the fluoroquinolones and may decrease the effectiveness of treatment with other quinolones due to the potential for cross resistance. Efficacy data have shown an insufficient efficacy of the product for the treatment of acute mastitis caused by Gram-positive strains.

Special precautions to be taken by the person administering the veterinary medicinal product to animals: None.

Pregnancy and lactation: Studies in laboratory animals (rats, rabbits) did not show any teratogenic, embryotoxic effects or any maternal toxicity of marbofloxacin. Safety

of the veterinary medicinal product at doses of 2 mg/kg has been shown in cows during gestation and in suckling pigs and calves when used in sows and cows. Safety of the veterinary medicinal product at doses of 8 mg/kg has not been shown in pregnant cows or in suckling calves when used in cows. Use only accordingly to the benefit/risk assessment by the responsible veterinarian. In the case of use in the cow during lactation, see withdrawal periods.

Interactions with other medicaments: None known.

Overdose: No signs of overdose have been observed after administration of 3 times the recommended dose. Signs such as acute neurological disorders may occur when the dose is exceeded. This signs should be treated symptomatically. Do not exceed the recommended dose.

Incompatibilities: In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

Disposal

Medicines should not be disposed of via wastewater or household waste.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required. These measures should help to protect the environment.

Other information

Properties: Marbofloxacin is a synthetic, bactericidal antimicrobial, belonging to the fluoroquinolone group, which acts by inhibition of DNA gyrase. It is effective in vitro against a wide range of Gram-positive bacteria (in particular *Staphylococcus*) and Gram-negative bacteria (*Escherichia coli*, *Pasteurella sp*) as well as *Mycoplasma* (*Mycoplasma bovis*). Resistance to *Streptococcus* may occur. After subcutaneous or intramuscular administration in cattle and pigs at the recommended dose of 2 mg/kg, marbofloxacin is readily absorbed and reaches maximal plasma concentrations of 1.5 µg/ml within less than 1 hour.

Its bioavailability is close to 100%. Marbofloxacin is weakly bound to plasma proteins (less than 10% in pigs and less than 30% in cattle) and extensively distributed. In most tissues (liver, kidney, skin, lung, bladder, uterus) it achieves higher concentrations than in plasma. After intramuscular administration in dairy cows, marbofloxacin reaches maximal milk concentration of 1.02 µg/ml (C_{max} after first administration) at 2.5 hours (T_{max} after first administration). Marbofloxacin is eliminated slowly in pre-ruminating calves (t_{1/2} = 5-9 hours) and pigs (t_{1/2} = 8-10 hours), faster in ruminant cattle (t_{1/2} = 4-7 hours) and predominantly in the active form in urine and faeces. After a single intramuscular administration in cattle at the recommended dose of 8 mg/kg, the maximum plasma concentration of marbofloxacin (C_{max}) is 7.3 µg/ml reached in = 0.78h (T_{max}). Binding to plasma proteins is about 30%. Marbofloxacin is eliminated slowly (t_{1/2} = 15.60 h), predominantly in the active form in urine and faeces.

Pack sizes: Vials of 250 ml. Vials are individually packed in a carton box. Six, ten or twelve vials are grouped as a clinical pack. Not all pack sizes may be marketed.

For animal treatment only. To be supplied only on veterinary prescription.

Name and address of the manufacturer:

Invesa
Industrial Veterinaria, S.A.
a LIVISTO company
Esmeralda, 19
08950 Esplugues de Llobregat
(Barcelona) Spain

Marketing authorisation holder:

YENHER AGRO PRODUCTS SDN. BHD. (222455-P)
1628 Jalan IKS Simpang Ampat 1,
Taman IKS Simpang Ampat.
14100 Simpang Ampat, S.P.S.,
Penang, Malaysia.

Date of revision of package insert: March 2019