

PREVIRON

Iron (III) (as gleptoferron) 200 mg/ml solution for injection

COMPOSITION PER ML:

Iron (III) 200 mg (as gleptoferron 532.6 mg); Phenol 5 mg;
Other excipients q.s.ad 1 ml.

PRODUCT DESCRIPTION:

Dark brown, slightly viscous solution.

PHARMACODYNAMICS/PHARMACOKINETICS:

PREVIRON is a veterinary medicinal product containing iron (III) as gleptoferron in solution for injection. Gleptoferron is an Iron-III-dextran glucoheptonic acid complex. Iron dextran complexes are widely used in veterinary therapeutics for the prevention of iron deficiency anaemia in piglets.

Iron is an essential micronutrient. It plays a major role in the oxygen transport of haemoglobin and myoglobin, as well as a key role in enzymes, such as cytochromes, catalases, and peroxidases. Iron has a high recovery rate from metabolism and food ingested. Thus, deficiency occurs only very rarely in adult animals.

After intramuscular injection, the iron complex is absorbed into the lymphatic tissue within 3 days. Here, the complex is split to release Fe^{3+} which is stored as ferritin in the main storage organs (e.g. liver, spleen and the reticuloendothelial system). In the blood, free Fe^{3+} binds to transferrin (transport form) and is mainly used for the synthesis of haemoglobin.

INDICATION:

For the prevention of iron deficiency anaemia in piglets.

DOSAGE, ROUTE AND METHOD OF ADMINISTRATION:

The product is administered as a single 1 ml (200 mg iron) dose by deep intramuscular injection. Inject once between the 1st and the 3rd day of life.

CONTRAINDICATIONS:

Do not use in piglets suspected to suffer from deficiency of vitamin E and/or selenium.

Do not use in case of hypersensitivity to the active substance.

Do not use in animals with diarrhoea.

Do not administer intravenously.

WARNINGS AND PRECAUTIONS:

Special warnings for each target species

None.

Special precautions for use in animals

It is advisable to stretch the skin at the injection site to minimise leakage after withdrawal of the needle.

Normal aseptic injection techniques should be practised.

Avoid the introduction of contamination during use.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Care should be taken to avoid accidental self-injection as well as mucous membrane contact, especially people with known hypersensitivity to iron dextran. In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

People with known hypersensitivity to the active substance (gleptoferron) or with hemochromatosis should avoid contact with the veterinary medicinal product.

Wash hands after use.

INTERACTION WITH OTHER MEDICAMENTS:

None.

STATEMENT ON USAGE DURING PREGNANCY AND LACTATION:

Not applicable.

ADVERSE EFFECTS/UNDESIRABLE EFFECTS:

Occasionally discolouration of the tissue and/or slight, soft swelling may be observed at the site of injection. This should disappear within a few days. Also hypersensitivity reactions can occur.

Rarely deaths have occurred in piglets following the administration of parenteral iron dextran preparations. These deaths have been associated with genetic factors or deficiency of vitamin E and/or selenium.

Very rarely piglets deaths have been reported which have been attributed to an increased susceptibility to infection due to temporary blocking of the reticuloendothelial system.

If you notice any serious effects or other effects not mentioned in this package leaflet, please inform your veterinary surgeon.

OVERDOSE AND TREATMENT:

Large amounts of iron administered by the parenteral route may result in transient reduced capacity of the immune system due to iron overload of lymph macrophages. Pain, inflammation reactions, abscess formation as well as persistent discolouration of muscle tissue at the injection site may occur. Poisoning arising from treatment may result in the following signs: pale mucous membranes, haemorrhagic gastroenteritis, vomiting, rapid heart rate, hypotension, breathlessness, oedema of the limbs, lameness, shock, death, liver damage. Supportive measures such as chelating agents can be used. Iron overdose can result in gastrointestinal signs such as diarrhoea or constipation. Treat symptomatically.

INCOMPATIBILITIES:

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

WITHDRAWAL PERIOD:

Meat and offal: Zero days.

SPECIAL STORAGE PRECAUTIONS:

Keep medicine out of reach of children.

Jauhi Ubat Dari Kanak-kanak.

Store below 30 °C (room temperature).

Discard the product within 28 days of opening.

DISPOSAL OF CONTAINERS:

Dispose waste material by incineration.

DATE OF REVISION OF PACKAGE INSERT:

1st of January 2026

PACK SIZES:

Box with 1 vial of 100 ml.

Box with 1 vial of 250 ml.

Registration holder:

HIPRA MALAYSIA SDN BHD

LOT NO. A-03-03A, LEVEL 3,

PLAZA MONT'KIARA, NO. 2,

JALAN KIARA, MONT'KIARA,

50480 KUALA LUMPUR,

WILAYAH PERSEKUTUAN KUALA LUMPUR

Reg. n° MAL19076005HA

Controlled Medicine. Ubat Terkawal.
FOR ANIMAL USE ONLY

HIPRA

LABORATORIOS HIPRA, S.A.

Avda. la Selva, 135 · 17170 Amer (Girona) SPAIN

Tel. (34) 972 43 06 60. Fax (34) 972 43 06 61. hipra@hipra.com

717477-20.0

02-26

717477-20.0