

Pharmasin® 250 mg/g Premix for Medicated Feeding Stuff

Keep out of reach of children.

STATEMENT OF ACTIVE SUBSTANCE:

Tylosin (as tylosin phosphate):
250 mg per g. (equivalent to
250 000 IU)

PRODUCT DESCRIPTION:

Pharmasin ® 250 mg/g Premix for
Medicated Feeding Stuff for Pigs,
Broilers and Pullets represents off-
white to beige granules.

PHARMACODYNAMICS:

Tylosin is a macrolide antibiotic
produced by a strain of
Streptomyces fradiae. It exerts its
antimicrobial effect by inhibiting
protein synthesis of susceptible
micro-organisms.

The tylosin spectrum of activity
includes Gram-positive bacteria,
some Gram – negative strains such
as *Pasteurella*, and *Mycoplasma*
spp. at concentrations of 16 µg/ml or
less.

PHARMACOKINETICS:

In most species peak plasma
concentrations have been attained 1
to 2 hours after administration of
tylosin. Compared to plasma levels
clearly higher tissue concentrations
have been observed. Tylosin was
extensively metabolized. Most of the
residues are excreted in faeces
predominantly consisting of tylosin A,
tylosin factor D and
dihydrodesmycosin.

INDICATIONS FOR USE:

Pigs: Treatment and prevention of
Porcine Intestinal Adenomatosis
(Ileitis) associated with *Lawsonia*
intracellularis when the disease has
been diagnosed at the group or herd
level.

Broilers and pullets: Treatment and
prevention of respiratory infections
caused by *Mycoplasma*
gallisepticum and *Mycoplasma*
synoviae, when the disease has
been diagnosed in the flock.
Treatment and prevention of necrotic
enteritis caused by *Clostridium*
perfringens, when the disease has
been diagnosed in the flock.

RECOMMENDED DOSAGE AND MODE OF ADMINISTRATION:

Administration through the feed: for
the preparation of a medicated feed
containing 40-1100 gram tylosine per
ton of feed, the required amount of
product should be homogenously
mixed with a suitable carrier into a
feed premixture so that at least 5 kg
of this premixture can be added to
the feed in order to obtain a
medicated feed with the required
concentration.

For the preparation of medicated feed:
As 1 kg of product contains 250 g
tylosin activity it follows that 4 mg
Pharmasin 250 mg/g premix
corresponds to 1 mg tylosin activity.
The dosages are as follows:

Pigs

For the treatment and prevention of
porcine intestinal adenomatosis (PIA):
4 – 5 mg tylosin per kg BW
(corresponding to 16 - 20 mg product
per kg BW) for 3 weeks.

Broilers and pullets:

For the treatment and prevention of
respiratory infections:

127 mg tylosin per kg BW
(corresponding to 508 mg product per
kg BW) for the first 5 days of life. It is
strongly recommended to repeat the
treatment of the birds at the age of
3-4 weeks.

For the treatment and prevention of
necrotic enteritis:

10 – 20 mg tylosin per kg BW
(corresponding to 40 – 80 mg product
per kg BW) for 7 days.

For the preparation of the medicated
feed the body weight of the animals to
be treated and their actual daily feed
consumption should be taken into due
account. Consumption may vary
depending on factors like age, breed,
husbandry system. To provide the
required amount of active substance
in mg per kg mixed feed the following
calculation should be made:

$$\frac{\text{.... mg product} \times \text{average body weight (kg) of}}{\text{/kg BW/day} \quad \text{the animals to be treated}} = \text{... mg product}$$

Average daily amount of mixed feed intake /kg per animal per kg/ mixed feed

The mixing should be performed by
an (authorised) feeding stuff
manufacturer with adequate mixing
apparatus.

The uptake of medicated feed
depends on the clinical condition of
the animals. In order to obtain the
correct dosage the concentration of
tylosin should be adjusted accordingly.
Should there be no clear response to
treatment within 3 days the treatment
approach should be reconsidered.
Body weight should be evaluated
accurately to avoid under dosing.

CONTRAINDICATIONS:

Do not use in animals with known
sensitivity to the active substance
and/or to any of the excipients of the
veterinary medicinal product.

Do not use in animals with known
hyper sensitivity to tylosin and other
macrolides.

Do not use where cross-resistance to
other macrolides (MLS-resistance) is
suspected.

Do not use in animals vaccinated with
tylosin-sensitive vaccines either at the
same time or within 1 week
previously.

Do not use in animals with hepatic
disorders.

Do not use in horses. Danger of
inflammation of the cecum.

WARNINGS AND PRECAUTIONS:

Special precautions for use in animals

Animals with acute infections may
have a reduced feed intake and
should be treated with a suitable
injectable product first.

Due to likely variability (time,
geographical) in susceptibility of
bacteria for tylosin, bacteriological
sampling and susceptibility testing are
recommended.

Inappropriate use of the veterinary
medicinal product may increase the
prevalence of bacteria resistant to
Tylosin and other macrolides.

Special precautions for the person administering the veterinary medicinal product to animals

Tylosin may induce irritation.
Macrolides, such a tylosin, may also
cause hypersensitivity (allergy)
following injection, inhalation,
ingestion or contact with skin or eye.
Hypersensitivity to tylosin may lead to
cross reactions to other macrolides
and vice versa. Allergic reactions to
these substances may occasionally
be serious and therefore direct
contact should be avoided.

STATEMENT ON USAGE DURING PREGNANCY AND LACTATION:

Laboratory studies in mice and rats
have not produced any evidence of
teratogenic, foetotoxic or
maternotoxic effects. No studies have
been conducted in the target species
population. Use only according to the
benefit/risk assessment by the
responsible veterinarian.

ADVERSE EFFECTS/

UNDESIRABLE EFFECTS:

In pigs, adverse reactions have been
observed, including diarrhoea,
pruritus, erythema, rectal oedema and
prolapse.

OVERDOSE AND TREATMENT:

Tylosin has been shown to produce
no adverse effects when fed to pigs at
600 ppm in the feed (three to six
times the recommended dose level)
for 28 days. At high levels diarrhoea,
apathy, convulsions may occur. The
therapy is symptomatic.

INCOMPATIBILITIES:

In the absence of compatibility
studies, this veterinary medicinal
product must not be mixed with other
veterinary medicinal products.

WITHDRAWAL PERIODS:

Meat & offal:

Pig: Zero days
Broilers and pullets: 1 day
Do not use in laying hens producing
eggs for human consumption.

DISPOSAL

Any unused veterinary medicinal
product or waste materials derived
from such veterinary medicinal
products should be disposed of in
accordance with local requirements.

STORAGE CONDITIONS:

Store in the original container to
protect from light. Store in a dry
place. Store below 30°C.

PACKAGING AVAILABLE:

20 kg paper/polyethylene bag.
1 kg resealable block bottom zipped
sachet made of polyethylene
/aluminium/polyethylene
terephthalate

MANUFACTURER:

BIOVET Joint Stock Company
39 Petar Rakov Street,
4550 Peshtera,
Bulgaria

MARKETING AUTHORIZATION HOLDER:

Pahang Pharmacy Sdn. Bhd.
(10826-K)
Lot 5979 Jalan Teratai, 5½ Mile Off
Jalan Meru, 41050 Klang,
Selangor Darul Ehsan,
Malaysia

DATE OF REVISION:

March, 2018.