

PELANGI VET PHARMA SDN. BHD.

No 8, PERDA Industrial Park, Lorong IKS, Simpang Ampat B,
14100 Simpang Ampat, S.P.S, Pulau Pinang, Malaysia

PRODUCT LITERATURE

PVP-LEVAMISOLE 20% W/V SOLUTION

Description:

Colourless clear solution

Each 1 L solution contains:

Levamisole HCl	262.00 g
----------------	----------

Preservatives:

- | | |
|-------------------|-----------|
| 1. Methyl Paraben | 2.00 g |
| 2. Propyl Paraben | 200.00 mg |

Pharmacodynamics:

Levamisole is the laevo isomer of tetramisole, a racemic imidazothiazole derivative, and is a member of the imidazothiazole group of anthelmintics.

Levamisole is a broad spectrum anthelmintic which displays excellent activity against mature and developing immature stages of gastro-intestinal and pulmonary nematodes.

By behaving as a cholinergic agonist of the nematode nervous system levamisole mimics the action of the excitatory neurotransmitter, acetylcholine, which results in sustained (spastic) muscle paralysis. By inhibiting fumarate reductase, levamisole also has a minor role to play in disrupting the nematodes energy pathway. However this is of limited consequence in comparison to its role as a paralysing agent.

Pharmacokinetics:

Levamisole is absorbed through the skin after dermal application. It is reportedly distributed throughout the body. Levamisole is primarily metabolized with less than 6% excreted unchanged in the urine. Plasma elimination half-lives have been determined for several veterinary species :

Cattle, 4-6 hours, Dogs, 1.8 – 4 hours and swine, 3.5 – 6.8 hours. Metabolites are excreted in both the urine (primarily) and feces

Indication:

Levamisole is a broad- spectrum anthelmintic indicated for use in cattle in the treatment and control of nematode infections such as parasitic gastro-enteritis and lungworm disease caused by the following mature and developing immature gastro-intestinal and pulmonary nematodes :

Lungworms - *Dictyocaulus viviparus*

Gastrointestinal worms - *Trichostrongylus* spp, *Cooperia* spp, *Ostertagia ostertagi* (except inhibited *O. ostertagi* larvae), *Haemonchus* spp, *Nematodirus* spp, *Bunostomum* spp, *Oesophagostomum* spp.

Warning & Precautions:

There are no known warning and precautions reported.

Drug Interactions:

Cattle must not be treated within a period of 14 days before or after treatment with organophosphorus.

Environmental Properties:

Any unused veterinary medical product or waste materials derived from such veterinary medical products should be disposed of in accordance with local requirements. Harmful to fish and aquatic life. Do not contaminate ponds, waterways or ditches with the product or empty container.

Pregnancy & Lactation:

May be administered to pregnant or lactating animals but care should be taken when treating heavily pregnant animals or animals suffering stress from adverse weather conditions, poor nutrition, penning, handling etc.

Side effects:

The product is safe for use in cattle at the recommended dosage. In cases of overdosage (which only occurs at over five times the recommended dose rate), hyperaesthesia, tremor and occasionally diarrhoea may occur.

At the recommended dose rates animals should not show any adverse side effects. However, local skin irritation at the application site may be observed occasionally, characterised by subcutaneous oedema. Severe cases may show signs of epidermal flaking for which symptomatic treatment is recommended.

Mode of Administration

Topical

Recommended Dose:

The pour-on is indicated for external transcutaneous administration to cattle.

The recommended dose rate is 10 mg levamisole/kg bodyweight equivalent to 2.5 ml per 50 kg bodyweight.

Bodyweight	Dose
Up to 50 kg (1 cwt)	2.5 ml
51-100 kg (1-2 cwt)	5 ml
101-150 kg (2-3 cwt)	7.5 ml
151-200 kg (3-4 cwt)	10 ml
201-250 kg (4-5 cwt)	12.5 ml
251-300 kg (5-6 cwt)	15 ml
301-350 kg (6-7 cwt)	17.5 ml

Above 350 kg give a further 2.5 ml for each additional 50 kg bodyweight.

Dose all young cattle in early summer and if possible move to clean pasture. Cattle should be dosed on first signs of lungworm infection.

For external administration only. Pour on from the calibrated dispenser or using a standard dosing gun. Apply along the flattest part of the backline at the rate of 10 mg levamisole/kg bodyweight, equivalent to 2.5 ml per 50 kg bodyweight.

Symptoms & Treatment of Overdose:

In cases of overdosage (which only occurs at over five times the recommended dose rate), hyperaesthesia, tremor and occasionally diarrhoea may occur.

Storage:

Store below 30° C in dry place, protected from light.

Dosage Form:

Liquid in the form of solution

Packing:

Packed in 1L HDPE plastic container with cap

Disposal of Container:

To be disposed as per local authority requirement

Shelf life:

2 years from date of manufacture

Manufactured & Distributed by:

Pelangi Vet Pharma Sdn. Bhd. (665097K)

Lot 8, PERDA Industrial Park,
Lorong IKS, Simpang Ampat B
14100 Simpang Ampat,
S.P.S Pulau Pinang, Malaysia

Date:

26/12/2025