

PROZIL fort 1%w/v

Injectable solution

COMPOSITION

Each ml contains
Acepromazine.....10 mg
(equivalent to 13.55 mg acepromazine maleate)
Benzyl alcohol (as preservative).....0.01 ml
Exp. qs.1 ml

DESCRIPTION: The finished product is yellow transparent solution.

PHARMACODYNAMICS

Acepromazine is a phenothiazine. It is a central nervous system depressant with associated activity on the autonomic system. Phenothiazines have a central action due to inhibition of dopamine pathways, resulting in alteration of mood, reduction in fear and removal of learned or conditioned responses. Acepromazine possesses anti-emetic, hypothermic, vasodilatory (and therefore hypotensive) and anti-spasmodic properties.

PHARMACOKINETICS

The length of action of acepromazine appears to be prolonged and to be dose dependent.

INDICATION

Anaesthetic Premedication: Following acepromazine administration, the amount of anaesthetic necessary to induce anaesthesia is considerably reduced.

Tranquilisation: Acepromazine tranquilisation (ataraxy) involves a modification of temperament which is not associated with hypnosis, narcosis or marked sedation. This is achieved with low doses of acepromazine. At low doses, acepromazine reduces anxiety which is beneficial for use in horses prior to shoeing or transportation.

Sedation: At higher dose rates acepromazine is an effective sedative, as an adjunct to, or replacement for, physical restraint e.g. dentistry, handling and shoeing. The relaxant effects aid examination of the penis in horses and the treatment of tetanus and choke.

TARGET SPECIES: Horses.

RECOMMENDED DOSE

For intramuscular or intravenous injection. In case of intravenous injection, it is recommended the injection is made slowly.

0.03-0.10 mg acepromazine per kg body weight, equivalent to 0.15-0.5 ml product per 50 kg body weight.

Normally, single doses of acepromazine are administered. Long term use is not recommended. On the rare occasions that repeat dosing is required, the dosing interval should be 36-48 hours.

Take adequate precautions to maintain sterility. Avoid the introduction of contamination during use. Should any apparent growth or

discolouration occur, discard the product.

ROUTE OF ADMINISTRATION

For intramuscular or intravenous injection. In case of intravenous injection, it is recommended the injection is made slowly.

CONTRA-INDICATION

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

Do not administer to breeding stallions..

Do not use in pregnant mares.

Do not use in animals in existing severe emotional excitation.

INTERACTION WITH OTHER MEDICAL PRODUCT

Phenothiazines are additive to the actions of other CNS depressants and will potentiate general anaesthesia (see indications for use).

Do not use this product in conjunction with organophosphates and/or procaine hydrochloride, as it may enhance activity and potential toxicity.

WARNINGS AND PRECAUTIONS

Acepromazine has little, if any, analgesic effect so that painful procedures must be avoided, particularly where animals are known to have unpredictable temperaments. Therefore, the usual precautions should be maintained when handling sedated horses.

During sedation, horses will normally retain visual and auditory acuity, so that loud sounds and rapid movements may cause arousal from the sedated state. It is therefore important to keep treated horses in a quiet environment and avoid sensory stimulation as far as possible.

Special precautions for use in animals

Situations may arise where general anaesthesia is required in the 4-6 hours following use of the product. In such cases care should be taken to reduce the dose of other premedicants and anaesthetic agents, particularly parenteral barbiturates, so as to avoid potentiation and additive depressant effects.

When administered to male horses (geldings or non-breeding stallions), use the lowest dose recommended to produce the required effect.

Acepromazine may cause hypothermia due to depression of the thermoregulatory centre and peripheral vasodilation.

Acepromazine is an adrenoceptor blocking drug and this causes hypotension and lowered haematocrit. The product should therefore be administered with great caution,

and at low dose rates only to debilitated horses and animals in states of hypovolaemia, anaemia and shock, or with cardiovascular disease. Rehydration should precede acepromazine administration. Duration of action may be prolonged and this should be remembered when riding, as acepromazine may affect performance and appear in drug tests for some time.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

This product contains a potent sedative; care should be taken, when handling and administering the product, to avoid accidental self-exposure.

In case of accidental self-injection, seek medical advice immediately and show the package insert or the label to the physician but DO NOT DRIVE as sedation and changes in blood pressure may occur. Symptomatic treatment may be required.

If accidental eye contamination occurs, flush gently with fresh running water for 15 minutes. Medical advice should be sought if irritation persists.

In the event of accidental skin contamination, contaminated clothing should be removed and the area washed with large amounts of soap and water. Medical advice should be sought if irritation persists.

Wash hands and exposed skin thoroughly after use.

- The product is not authorised for use in horses intended for human consumption.

- Keep out of reach of children

- Jauhkan daripada kanak-kanak

PREGNANCY AND LACTATION

Do not administer to pregnant mares. The safety of the veterinary medicinal product has not been established during pregnancy or lactation.

SIDE EFFECTS

Acepromazine may cause hypotension (common) and lowered haematocrit (very common). Reversible paralysis of the retractor penis muscle has been associated with the use of parenterally administered acepromazine in horses (very common). Acepromazine has caused paraphimosis (uncommon), sometimes as a sequel to priapism, however this only very rarely results in permanent penile dysfunction. When extrusion of the penis occurs, the owner should be advised to inform his veterinary surgeon if retraction of the penis does not take place within 2-3 hours. Suitable treatments have been described in the veterinary literature e.g. manual compression during the period of general anaesthesia, penile support

and manual compression, use of an Esmarch bandage, or drug reversal (e.g. slow intravenous administration of benztrapine mesylate).

Accidental intracarotid injection in horses can produce clinical signs ranging from disorientation to convulsive seizures and death.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reactions(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports)

SYMPTOMS AND TREATMENT OF OVERDOSE

Transient dose-dependent hypotension may occur in cases of accidental overdose. Therapy should consist of discontinuing any other hypotensive treatment, supportive care such as intravenous infusion of warm isotonic saline to correct hypotension and close monitoring. In severe cases treatment with norepinephrine may be indicated but its use must be based on a careful evaluation of the benefit risk balance by the responsible veterinary surgeon.

Epinephrine (adrenaline) is contra-indicated in the treatment of acute hypotension produced by overdosage of acepromazine maleate, since further depression of systemic blood pressure can result.

INCOMPATIBILITIES

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

STORAGE CONDITION

Store at temperature not exceeding 30°C.

Strictly keep away from sunlight and avoid heat

SHELF LIFE: 2 years

SHELF LIFE AFTER OPENING: 24 hours

PACKAGING: 100 ml

REVISED DATE: Jun 24, 2019



Manufactured by: Vemedim Animal Health
Song Hau Industrial Zone, Chau Thanh District,
Hau Giang Province, Vietnam

Distributed by: Vemedim SDN. BHD.
15, Jalan USJ 1/31, 47600 Subang Jaya Selangor,
Darul Ehsan, Malaysia
Tel: (603)80247118 Fax: (603) 80247168