

Cocci-nil 45 mg/ml Oral Solution

(FOR VETERINARY USE ONLY)

COMPOSITION

Each 1000 ml contents:

Sulfaquinoxaline45g
Benzyl alcohol (as preservative) 18.5 ml

PRODUCT DESCRIPTION

The product is a yellowish colour oral solution.

PHARMACODYNAMICS

Sulfonamides are usually bacteriostatic agents when used alone.

Sulfonamides interfere with the biosynthesis of folic acid in bacterial cells; they compete with para-aminobenzoic acid (PABA) for incorporation in the folic acid molecule. By replacing the PABA molecule and preventing the folic acid formation required for DNA synthesis, the sulfonamides prevent multiplication of the bacterial cell. Only organisms that synthesize their own folic acid are susceptible; mammalian cells use preformed folic acid and, therefore, are not susceptible. Cells that produce excess PABA or environments with PABA, such as necrotic tissues, allow for resistance by competition with the sulfonamide.

PHARMACOKINETICS

Most sulfonamides are well absorbed orally with the exception of the enteric sulfonamides, such as sulfaquinoxaline, which are minimally absorbed. Delays in absorption may occur in adult ruminants or when sulfonamides are administered with food to monogastric animals.

Sulfonamides are widely distributed throughout the body. They cross the placenta, and a few penetrate into the cerebrospinal fluid.

Sulfonamides are primarily metabolized in the liver but metabolism also occurs in other tissues. Biotransformation occurs mainly by acetylation, glucuronide conjugation, and aromatic hydroxylation in many species. The types of metabolites formed and the amount of each varies depending on the specific sulfonamide administered; the species, age, diet, and environment of the animal; the presence of disease; and, with the exception of pigs and ruminants, even the sex of the animal.

INDICATIONS

Chickens:

As an aid in the control of acute fowl cholera caused by *Pasteurella multocida* susceptible to sulfaquinoxaline and fowl typhoid caused by *Salmonella gallinarum* susceptible to sulfaquinoxaline.

As an aid in the control of outbreaks of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, and *E. brunetti*.

CONTRA-INDICATIONS

Hypersensitive to sulfas, severe renal or hepatic impairment.

RECOMMENDED DOSE:

Oral Administration through drinking water

Prepare medication drinking water fresh daily

Chickens

For control of acute fowl cholera and fowl typhoid:

Administer at the 0.04 percent level in drinking water (approximately 4.45 L product/ 500 L drinking water) for 2 or 3 days. Move birds to clean ground. If disease recurs, repeat treatment. If cholera has become established as the respiratory or chronic form, use feed medicated with sulfaquinoxaline. Poultry which have survived typhoid outbreaks should not be kept for laying house replacements or breeders unless tests show they are not carriers.

For control of outbreaks of coccidiosis:

Administer at the 0.04 percent level in drinking water (approximately 4.45 L product/ 500 L drinking water) for 2 or 3 days, skip 3 days then administer at the 0.025 percent level in drinking water (approximately 2.78 L product/ 500 L drinking water) for 2 more days. If bloody droppings appear, repeat treatment at the 0.025 percent level in drinking water (approximately 2.78 L product/ 500 L drinking water) for 2 more days.

Medicated chickens must actually consume enough medicated water which provides a recommended dosage of approximately 22 to 100 mg per kg per day in chickens depending on the age, class of animal, ambient temperature, and other factors.

WARNING AND PRECAUTIONS

- May cause toxic reactions unless the drug is evenly mixed in water at dosages indicated and used according to directions.
- Do not give flushing mashes.
- People who handle this medication should avoid contact with eyes, skin, or clothing to prevent eye and skin burns. In case of contact, the areas affected should be flushed for at least fifteen minutes; medical attention should be sought for eye exposure.
- Keep medicine out of the reach of children.
- Jauhkan ubat daripada kanak-kanak.

INTERACTION WITH OTHER MEDICAMENTS

Drug interaction relating specifically to the use of sulfonamides in animals are rarely reported in veterinary literature.

PREGNANCY AND LACTATION

This product must not be used on laying birds

SIDE EFFECT

Prolonged administration of sulfaquinoxaline may result in depressed feed intake, decomposition of crystals in the kidney, or interference with normal blood clotting. Hemorrhagic syndrome (anorexia, epistaxis, hemoptysis, lethargy, pale mucous membranes, possibly death) has been reported in chickens but may occur in other species. Sulfaquinoxaline is a vitamin K antagonist that inhibits vitamin K eposide and vitamin K quinone reductase and causes an effect similar to that of coumarin anticoagulants.

OVERDOSE AND TREATMENT

Toxicities secondary to acute overdose of sulfonamides are not typically reported. Side effects may be more likely to occur with high doses and long-term administration, but are seen at recommended doses as well.

WITHDRAWAL PERIODS

Do not give to chickens within 10 days of slaughter for food.
Do not medicate chickens producing eggs for human consumption.

SHELF LIFE

36 months from manufacturing date
After first opening : 24 hours
After reconstitution or dilution : 24 hours

STORAGE

Store in a cool and dry room ($\leq 30^{\circ}\text{C}$). Keep away from direct sunlight

MAL NO. :

PACKAGING : 1 L
EXP DATE :
MFG DATE :
Date of Revision : 29th July 2019

Manufacturer:

Shennong Animal Health (M) Sdn. Bhd.
6, 8, & 10, Jalan Industri USJ 1/19,
Taman Perindustrian USJ 1,
47610 Subang Jaya, Selangor, Malaysia.

Product registration holder:

F.E Venture Sdn. Bhd.
3 & 5, Jalan PJS 11/8, Bandar Sunway,
46150 Petaling Jaya, Selangor D.E., Malaysia