

MUCOSTOP 600 MG EFFERVESCENT TABLET

NAME AND STRENGTH OF ACTIVE SUBSTANCE

Each effervescent tablet contains 600mg Acetylcysteine

PRODUCT DESCRIPTION

White, round, flat-faced tablets.

Appearance of solution should be clear after dissolving in water.

PHARMACODYNAMICS

Acetylcysteine has strong fluidification activity to mucus secretion and mucopurulent by depolymerization of mucoprotein acid complex and nucleic acid were as act as viscosity agent from mucoid component, sputum purulenta and others secretions, it acts as anti fiogistic and increase mucosa regeneration speed.

PHARMACOKINETICS

Absorption

Bioavailability

Absorbed following oral administration, with peak plasma concentration achieved within 1 – 2 hours.

Absorbed from the pulmonary ephitelium.

Distribution

Extent: Crosses the placenta following oral or IV administration

Plasma protein binding: 50 – 83%

Metabolism

Deacetylated in the liver to cysteine

Elimination

Elimination route: Principally (70%) nonrenal

Half – life: 6.25 hours after oral administration

INDICATIONS

All respiratory tract diseases leading to the formation of thick secretions difficult to be expectorated, such as acute and chronic bronchitis, laryngitis, sinusitis, tracheitis, influenza, bronchial asthma and (as complementary treatment) mucoviscidosis.

RECOMMENDED DOSAGE

Adults:

1 effervescent tablet a day (preferably in the evening).

Dissolve in water before administration.

ROUTE OF ADMINISTRATION: Oral; Dissolve the effervescent tablet in a glass of water before consuming.

CONTRA-INDICATIONS

Patients with hypersensitivity to Acetylcysteine.

Contraindicated in children below two (2) years of age.

As Mucostop 600mg Effervescent Tablet contains aspartame, it is contraindicated in patients suffering from phenylketonuria.

Active peptic ulcer.

WARNING AND PRECAUTIONS:

Unsuitable for phenylketonurics.

Patient with gastric, should be administered after food.

Several animal and human researches showed no teratogenic and dangerous adverse effect to pregnant and nursing patient, however administer only at several condition needs and always controlled by physician.

Patients suffering from bronchial asthma must be strictly controlled during the therapy; should bronchospasm occur, the treatment must immediately be suspended.

Caution should be taken in patient suffering from or with a history of peptic ulcer.

The possible presence of a sulphurous odour does not indicate an alteration of the product but is a characteristic of the active ingredient contained in this preparation.

It is preferable not to mix other drugs with the Mucostop 600 solution.

INTERACTIONS WITH OTHER MEDICAMENTS

No cases of drug-drug interaction are described in literature for NAC by oral use.

PREGNANCY AND LACTATION

Pregnancy: category B

Lactation: Not known whether acetylcysteine is distributed into milk; caution advised.

ADVERSE REACTIONS

Hypersensitivity reactions have been reported in patients receiving Acetylcysteine, including bronchospasm, angioedema, rashes and pruritus. Other adverse effects reported with Acetylcysteine include nausea and vomiting, fever, syncope, sweating, arthralgia, blurred vision, disturbances or liver function.

Immune system disorders: Anaphylactic / anaphylactoid reaction

Skin and subcutaneous Tissue Disorders: Severe cutaneous adverse reactions (SCAR) eg erythema multiforme, Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN). In most of these cases reported at least one other drug was administered at the same time, which may have possibly enhanced the described mucocutaneous effects.

OVERDOSE AND TREATMENT

In the cases of accidental or non-accidental overdose, it may cause irritation, pyrosis, stomachache, nausea, vomiting and diarrhea. There are no serious side effects or toxic symptoms reported, even in patients treated with very high doses of acetylcysteine.

However, it is advisable to consult the physician in any case of overdoses.

STORAGE

Store at room temperature below 30°C and dry place.

DOSAGE FORM AND PACKAGING AVAILABLE

10 effervescent tablets in 1 plastic tube and silica gel cap

Manufacturer

Neutec İlaç San. Tic. A.Ş.

1. OSB 1. Yol No: 3

Adapazarı / SAKARYA / TURKEY

Product Registration Holder

MEDISPEC (M) SDN BHD
55 & 57, Lorong Sempadan 2,
(Off Boundary Road)
11400 Ayer Itam, Penang,
Malaysia

Date of revision

14 October 2019