

Veterinary Package Insert

TP-FLUMIX 200 MG/ML SOLUTION

PRODUCT DESCRIPTION

Light yellow to yellow colour liquid, which contains flumequine 200mg in 1ml of solution.

PHARMACODYNAMICS & PHARMACOKINETICS

Flumequine is an antibacterial agent with bactericidal activity, belonging to the group of fluoroquinolones. It acts on the DNA, causing a blockade of the replication by inhibition of the A subunit of the DNA-gyrase enzyme of the bacterium, an enzyme involved in the formation of the DNA helix.

After oral administration, 10% of the dose is absorbed, reaching its maximum plasma concentration at 2 hours. It is well distributed in all tissues. The binding to plasma proteins is 74.5%. In blood is unchanged flumequine and its hydroxylated metabolite, which has less activity. In urine, 40-60% of the absorbed dose is excreted in the form of an unchanged drug, and the rest in the form of an inactive glucuroconjugate metabolite. A small portion is excreted in stool.

INDICATION

Poultry (broilers): Treatment of infections caused by bacteria sensitive to flumequine, such as *Pasteurella multocida* and *E. coli*.

Bovine (pre-ruminant calves): Treatment of infections caused by bacteria sensitive to flumequine, such as *Pasteurella multocida*, *E. coli* and *Salmonella* spp.

TARGET SPECIES

Poultry (broilers) and bovine (pre-ruminant calves).

RECOMMENDED DOSAGE AND ADMINISTRATION

Administration in drinking water.

Dosage for poultry (broilers): 18 mg flumequine / kg p.v./day for 5 days (equivalent to 0.09 ml of medicine / kg p.v./day) to add to the drinking water according to the consumption thereof.

Dosage for bovine (pre-ruminant calves): the total duration of the treatment is 5 days starting with a first dose of 12 mg of flumequine / kg p.v., (equivalent to 0.06 ml of medicine / kg p.v.) said dose is to be added to the drinking water according to the consumption thereof.

The first dose must be followed by 9 administrations of 6 mg of flumequine / kg pv / 12

hours (equivalent to 0.03 ml of medicine / kg pv / 12 hours) these doses are to be added to the drinking water according to the consumption of the same.

For poultry (broilers) and for bovine (pre-ruminant calves), the amount of medicine to be added to water (ml of medicine / liter of drinking water), can be calculated with the following formula:

$$\frac{\text{ml of medication}}{\text{liter of drinking water}} =$$

$$\frac{\text{dose of active substance (mg/kg p.v./ day)} \times \text{average weight of animals}}{\text{ml of active substance in ml of medicine (mg/ml)} \times \text{average water consumption (liter/day)}}$$

$$\frac{\text{dose of active substance (mg/kg p.v./ day)} \times \text{average weight of animals}}{\text{ml of active substance in ml of medicine (mg/ml)} \times \text{average water consumption (liter/day)}}$$

CONTRAINDICATIONS

Do not use in animals with renal or hepatic insufficiency.

WARNINGS & PRECAUTIONS.

i.) Special warnings for each target species

It is not allowed to be used in laying eggs which are for human consumption.

ii.) Special precautions for use

This medicine does not contain any antimicrobial preservative.

Acute cases and seriously ill animals with a reduction in water intake should be treated parenterally.

iii.) Special precautions for use in animals

Administer in the drinking water, ensuring a homogeneous mixture.

Do not expose treated animals to the sun, due to the risk of photosensitization.

When using this product should take into account the official recommendations on the use of antimicrobials.

The use of fluoroquinolones should be reserved for the treatment of those clinical cases that have responded poorly, or are expected to respond poorly, to other classes of antimicrobials.

The use of the product under conditions other than those recommended in the Technical Data Sheet may increase the prevalence of bacteria resistant to fluoroquinolones and decrease the effectiveness of treatment with other quinolones due to cross-resistance.

Avoid co-administration with foods containing divalent and trivalent cations or leave the product in metal containers.

iv.) Specific precautions to be taken by the person administering the veterinary medicinal product to animals

People with known hypersensitivity to fluoroquinolones should not manipulate the product.

During the preparation of medicated water, wash any splashes in eyes and skin with water. Wash hands after use.

Do not eat, drink or smoke while handling the medication.

INTERACTIONS WITH OTHER MEDICATIONS

Do not associate with trimethoprim, theophylline. The active substance interacts with divalent and trivalent cations.

PREGNANCY AND LACTATION

No contraindication has been described during these periods.

ADVERSE EFFECTS

On very rare occasions it can occur:

- Digestive disorders such as vomiting and diarrhea.
- Nervous disorders such as hyperexcitability of the animal.
- Phenomena of photosensitivity and cutaneous manifestations of allergic type.
- In case of prolonged use, alteration of joint development in young animals.

OVERDOSE & TREATMENT

In case of overdose may appear digestive or nervous alterations, which remit when stopping the medication.

WITHDRAWAL PERIOD

Meat: Broilers: 2 days.

Calves (pre-ruminants): 6 days.

MAXIMUM RESIDUAL LIMIT (MRL)

Substance / Drug Definition of residues in which MRL was set	Food	Maximum Residue Limits (MRLs) µg/kg
Flumequine	Muscle, fat	50
	Liver	100
	Kidney	300

STORAGE CONDITION

Store below 30 °C. Protect from direct sunlight.

SHELF LIFE

3 years.

After 1st opening, reconstitution or dilution, use within 24 hours.

PACKING

1 L

THYE PHARMA SDN. BHD.

(manufacturer & product registration holder)

No.2 Lorong Industri Ringan Permatang

Tinggi 8,

Kawasan Industri Ringan Permatang Tinggi,

14000 Bukit Mertajam,

Pulau Pinang.

Revised August 2019