

ENROTRON FLAVOUR 50 mg

Tablets Enrofloxacin

Composition

1 tablet contains: Enrofloxacin, 50.0 mg.
White to off-white, round tablet with one break line and one decorative line. The tablet can be divided into two equal parts.

Indications

For the treatment of bacterial single or combined infections of the respiratory, alimentary or urinary tract, the skin or wounds, caused by enrofloxacin-sensitive Gram-negative and Gram-positive bacteria: *E. coli*, *Pasteurella* spp., *Haemophilus* spp. and staphylococci.

Contraindications

Do not use in young or growing dogs (dogs aged less than 12 months (small breed) or less than 18 months (large breed)) as the product may cause epiphyseal cartilage alterations in growing puppies.
Do not use in dogs having seizure disorders, since enrofloxacin may cause CNS stimulation.
Do not use in dogs with known hypersensitivity to fluoroquinolones or to any of the excipients of the product.
Do not use in case of resistance to quinolones, as there exists almost complete cross resistance to other quinolones and complete cross resistance to other fluoroquinolones.
Do not use with tetracyclines, phenicols or macrolides because of potential antagonistic effects.
See also section "Special warnings".

Adverse reactions

In rare cases, vomiting and diarrhoea are observed.
See section "Contraindications".
If you notice any serious effects or other effects not mentioned in this package leaflet, please inform your veterinary surgeon.

Target species

Dogs.

Dosage for each species, route and method of administration

Tablet for oral use.
5 mg enrofloxacin per kg bw daily, corresponding to ½ tablet per 5 kg b.w. and 1 tablet per 10 kg b.w.
Can be administered directly or given with food.
Treatment generally takes place over 5 – 10 consecutive days.
If there is no clinical improvement within 3 days, the sensitivity test is to be repeated and it may be necessary to switch to a different treatment.

The recommended doses should not be exceeded.
To ensure a correct dosage, body weight should be determined as accurately as possible to avoid underdosing.

Withdrawal period

Not applicable.

Storage precautions

Keep out of the sight and reach of children.
Store in the original package in order to protect from light.
Divided tablets should be returned in the original package.
Any divided tablet portions remaining after 72 hours should be discarded.
Do not store above 30 °C.

Special warnings

Special warnings for each target species: None.

Special precautions for use in animals: Official and local antimicrobial policies should be taken into account when the product is used. Fluoroquinolones should be reserved for the treatment of clinical conditions which have responded poorly, or are expected to respond poorly, to other classes of antimicrobials.

Whenever possible, fluoroquinolones should only be used based on susceptibility testing. Use of the product deviating from the instructions given in the package insert may increase the prevalence of bacteria resistant to fluoroquinolones and may decrease the effectiveness of treatment with other quinolones due to the potential cross resistance.

As enrofloxacin is metabolized by the liver and partly eliminated via the kidneys, elimination may be delayed in dogs with liver or renal disturbances. Therefore, in cases of known liver or renal impairment, the product should be used with caution.

Special precautions to be taken by the person administering the veterinary medicinal product to animals: People with known hypersensitivity to fluoroquinolones should avoid contact with the product. Wash hands after use. In case of contact with the eyes, wash immediately with plenty of clean water. In case of accidental ingestion, seek medical advice immediately and show the package leaflet to the physician.

Pregnancy and lactation: Do not use during pregnancy and lactation.

Interactions: Concurrent use of flunixin should be under careful veterinary monitoring, as the interactions between these drugs may lead to adverse events related to delayed elimination. Elimination of theophylline may be delayed. Antagonistic effect may occur if enrofloxacin is combined

with phenicols, macrolide antibiotics or tetracyclines. Substances containing magnesium or aluminium, if administered at the same time, may impair the enrofloxacin re-sorption. Fluoroquinolones are fully cross-resistant.

Overdose: Overdosing can cause vomiting and nervous signs (muscle tremor, incoordination and convulsions) which may require treatment discontinuation.

In the absence of any known antidote, apply drug elimination methods and symptomatic treatment. If necessary, administration of aluminium- or magnesium-containing antacids or activated carbon can be used to reduce absorption of enrofloxacin.

According to literature, signs of overdose with enrofloxacin in dogs such as inappetence and gastrointestinal disturbance were observed at approximately 10 times the recommended dose when administered for two weeks. No signs of intolerance were observed in dogs administered 5 times the recommended dose for a month.

Disposal

Medicines should not be disposed of via wastewater or household waste.

Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

Other information

Enrofloxacin is a synthetic fluoroquinolone antibiotic that exerts its activity by inhibiting topoisomerase II, an enzyme involved in the mechanism of bacterial replication. Enrofloxacin exerts bactericidal activity concentration-dependent with similar values of minimal inhibit concentration and minimal bactericide concentrations. It also possesses activity against bacteria in the stationary phase by an alteration of the permeability of the outer membrane phospholipid cell wall.

Enrofloxacin is active against a wide range of Gram-negative bacteria, against mycoplasmas and against many Gram-positive bacteria.

Pseudomonas aeruginosa is variably susceptible and, when it is susceptible, usually has a higher MIC than other susceptible organisms.

Induction of resistance against quinolones can develop by mutations in the gyrase gene of bacteria and by changes in cell permeability towards quinolones.

Enrofloxacin is rapidly metabolised to form an active compound, ciprofloxacin.

After oral administration of Enrotron Flavour 150 mg (5 mg/kg) in dogs:

The maximal plasma concentration of enrofloxacin of 1205.48 ng/ml was observed 1.42 hours following administration.

The maximal plasma concentration of ciprofloxacin (432.48 ng/ml) was observed 2.6 hours following administration.

Enrofloxacin is primarily excreted via the kidneys. A major portion of the parent drug and its metabolites is recovered in urine.

Enrofloxacin is widely distributed in the body. The tissue concentrations are often higher than the serum concentrations. Enrofloxacin crosses the blood-brain barrier.

The degree of protein binding in serum is 14 % in dogs. In dogs (5 mg/kg), the half-life for enrofloxacin in serum is 2.73 hours and the half-life for ciprofloxacin is 5.35 hours. Approximately 60 % of the dose is excreted as unchanged enrofloxacin and the remainder as metabolites, amongst others ciprofloxacin. The enrofloxacin clearance is approximately 28.2 ml/minute/kg bw in dogs and the ciprofloxacin clearance is approximately 30.8 ml/minute/kg bw in dogs.

Bioavailability is in excess of 80 %.

Pack sizes: Box with 50 or 100 tablets packed in aluminium/aluminium blister foils of 10 tablets.

For animal treatment only. To be supplied only on veterinary prescription.

Name and address of the manufacturer

LelyPharma B. V.
Zuiveringweg 42
8243 PZ Lelystad
The Netherlands

Marketing authorisation holder

My Vet Care Sdn. Bhd.
A-G-3A, Red Ruby Shop Apartment
Jalan Indah 2/4, Taman Universiti Indah
43300 Seri Kembangan
Selangor, Malaysia

Date of revision of package insert: September 2019