

Hemaron Tablet

Hesperidin (Micronised) 50mg and Diosmin (Micronised) 450mg

Brand or Product Name

Hemaron Tablet

Name and Strength of Active Substance(s)

Hesperidin (Micronised) 50mg

Diosmin (Micronised) 450mg

Product Description

A brown oblong shaped film coated tablet.

Pharmacodynamics

Venotonic and vascular protector.

Pharmacology

It is active upon the return vascular system in the following way:

- It reduces venous distensibility and stasis,
- In the microcirculation, it normalises capillary permeability and increases capillary resistance.

Pharmacokinetics

In humans, following oral administration of Diosmin:

- Excretion is essentially faecal and urinary excretion is on average 14% of the administered quantity,
- The elimination half-life is of 11 hours.
- The product is highly metabolised, this metabolism is revealed by the presence of different phenol acids in the urine.

Indication

- Treatment of symptoms related to venolymphatic insufficiency (heavy legs, pain, early morning restless legs).
- Treatment of functional symptoms related to acute hemorrhoidal attack.

Dose/ Use Instruction

- Usual dosage: 2 tablets daily in two divided doses, midday and evening at meal times.
- Acute hemorrhoidal attack: 6 tablets per day for the first 4 days, then 4 tablets per day for 3 days.

Contraindications

Hypersensitivity to any of the components of Hemaron Tablet.

Warnings and Precautions

Acute hemorrhoidal attack: Administration of this medicine is no substitute for the specific treatment of other anal disorders. The treatment must be short-term. If the symptoms do not disappear rapidly, proctological examination should be performed and the treatment reviewed.

Interactions with Other Medications

Not Applicable

Pregnancy and lactation

No adverse effects have been reported to date in humans.

In the absence of data concerning the diffusion into breast milk, breastfeeding is not recommended during treatment.

Hemaron Tablet

Hesperidin (Micronised) 50mg and Diosmin (Micronised) 450mg

Adverse Effects/ Undesirable Effects

Nervous system disorders

Rare: dizziness, headache, malaise.

Gastrointestinal disorders

Common: diarrhea, dyspepsia, nausea, vomiting.

Uncommon: colitis.

Unknown frequency: abdominal pain.

Skin and subcutaneous tissue disorders

Rare: rash, pruritus, urticaria

Frequency unknown: isolated edema of the face, eyelids, lips. Exceptionally, angioedema.

Overdose and Treatment

Not Applicable

Storage Conditions

Store below 30°C.

Presentation:

PVC/Aluminium blister of 10 tablets.

Pack Size: 10, 30, 50, 100, 300, 500, and 1000 tablets

Manufactured by & Product Registration Holder:

Noripharma Sdn. Bhd. (792633-A)

Lot 5030, Jalan Teratai,

5 ½ Mile Off Jalan Meru,

41050 Klang, Selangor, Malaysia.

Date of Revision:

09th May 2019