

Euvichol-Plus Suspension Oral Cholera Vaccine
***Vibrio cholerae* O1 Inaba Cairo 48 (Heat inactivated)**
***Vibrio cholerae* O1 Inaba Phil 6973 El Tor (Formalin inactivated)**
***Vibrio cholerae* O1 Ogawa Cairo 50 (Formalin inactivated)**
***Vibrio cholerae* O1 Ogawa Cairo 50 (Heat inactivated)**
***Vibrio cholerae* O139 4260B (Formalin inactivated)**

Product Name

Euvichol-Plus Suspension Oral Cholera Vaccine

Product Description

A plastic tube filled with a yellow to yellowish suspension consists of inactivated *V. cholerae*

Composition

One dose (1.5 mL) contains:

Active ingredients:

V. cholerae O1 Inaba Cairo 48 classical biotype, Heat inactivated.....300 L.E.U
V. cholerae O1 Inaba Phil 6973 El Tor biotype, Formalin inactivated600 L.E.U
V. cholerae O1 Ogawa Cairo 50 classical biotype, Formalin inactivated.....300 L.E.U
V. cholerae O1 Ogawa Cairo 50 classical biotype, Heat inactivated.....300 L.E.U
V. cholerae O139 4260B, Formalin inactivated600 L.E.U

Indication

Euvichol-plus is indicated for active immunization against *Vibrio cholerae* but the efficacy against *Vibrio cholerae* serogroup O139 was not demonstrated.

The vaccine can be administered to anyone above the age of 1 year. Data for the safety and efficacy of the vaccine in infants (less than 1 year of age) is not available. The earliest onset of protection can be expected 7-10 days after the completion of the primary series of the vaccine.

Recommended Dosage

Two doses of vaccines should be given at an interval of two weeks.

Route of Administration

The vaccine is only recommended for oral administration.

Contraindications

1. The vaccine should not be administered to persons with either known hypersensitivity to any component of the vaccine, or having shown signs of severe reaction due to the previously taken dose.
2. Immunization with Euvichol-Plus should be delayed in the presence of any acute illness, including acute gastrointestinal illness or acute febrile illness.

Warnings and Precautions

1. As with any vaccine, immunization with Euvichol-Plus may not protect 100% of susceptible persons.
2. As with all vaccines, appropriate medical treatment should always be readily available in case of a rare event of anaphylactic reactions following the administration of the vaccine. For this reason, it is recommended that the person should remain under medical supervision for at least 30 minutes after vaccination.
3. This vaccine contains residual formaldehyde. Caution should be taken in subjects with known hypersensitivity to formaldehyde.
4. The safety and immune response of Euvichol-Plus has not been clinically evaluated in individual with HIV/AIDS.
5. No clinical study has been performed to evaluate the efficacy and safety of Euvichol-Plus in infants (less than 1 year of age). Therefore, the vaccine is not recommended for use in infants.

Interactions with Other Medicaments

Appropriate studies have not been conducted to determine the interaction with other medicinal products.

Pregnancy and Lactation

No specific clinical studies have been conducted to evaluate the efficacy and safety of Euvichol-Plus in pregnant or lactating women and for the fetus. Therefore, the vaccine is not recommended for use in pregnancy. Nevertheless, if you are exposed to or will be travelling to an area at risk for cholera, carefully evaluate the benefits and risks of administering this vaccine before considering it.

Side Effects

2,999 healthy children and adults (1-40 years) were participated in the clinical study for evaluating the safety.

- After taking the vaccines, during first 7 days, the most frequently reported adverse drug reactions in the clinical trial were headache, fever, diarrhea, nausea/vomiting and myalgia and 102 subjects (3.40%) among 2,999 subjects were reported. The incidence rate for children and adults is described on the table below.

	Total (N=2,999)	1 ~ 17 years (N=1,118)	18~40 years (N=1,881)
Total	3.40%	3.04%	3.62%
Headache	1.83%	0.81%	2.45%
Fever	1.00%	1.97%	0.43%
Diarrhea	0.67%	0.54%	0.74%
Nausea/Vomiting	0.37%	0.63%	0.21%
Myalgia	0.10%	0.00%	0.16%

- After taking the vaccines, adverse drug reactions were examined for a period of 28 days. 69 subjects (2.30%) among 2,999 subjects were reported with the adverse effects, and gastrointestinal disorders were reported the highest numbers i.e., 35 subjects (1.17%). The adverse drug reactions during the study (28 days) were described on the table below.
(Uncommon: 0.1~5%, Rare: less than 0.1%)

	Incidence rate	
	Uncommon	Rare
Gastrointestinal disorders	Abdominal pain, toothache, Diarrhea	Vomiting, abdominal pain upper
General disorders and administration site condition	Pyrexia	Thirst
Infection and infestations	Nasopharyngitis	Gastroenteritis
Nervous system disorders	Headache	Dizziness
Respiratory, thoracic and mediastinal disorders	Cough	Oropharyngeal pain
Skin and subcutaneous tissue disorders	Pruritus	Rash macular
Musculoskeletal and connective tissue disorders	-	Arthralgia, neck pain, pain in extremity
Vascular disorders	-	Flushing

Symptoms and Treatment of Overdose

No case of overdose has been reported.

Instructions for Use

1. The vaccine should be administered to anyone above the age of 1 year.
2. Two doses of vaccine should be given at an interval of two weeks.
3. The vaccine is presented as a suspension. Therefore, after shaking the vaccine container rigorously, 1.5 mL of the vaccine should be squirted into the mouth. Take a sip of water if necessary.
4. The frozen vaccines should not be taken.
5. The vaccine should not be administered parenterally (intramuscularly, subcutaneously or intravenously).

Presentation


1.5 mL of the final formulated bulk of Euvichol-Plus is filled into a 2 mL of plastic tube (LDPE).

1 strip is composed with 5 doses of plastic tube.

One folded paper box contains 1 strip (5 doses) or 2 strips (10 doses) or 10 strips (50 doses).

Storage Condition

The vaccine should be stored at 2°C ~ 8°C. Do not freeze.



Vaccine Vial Monitors (VVMs) are a part of the label attached on Euvichol-Plus supplied through EuBiologics Co., Ltd. The color dot which appears on the label of the vaccine is a VVM. This is a time-temperature sensitive dot that provides an indication of the cumulative heat to which the vaccine has been exposed. It warns the end user when exposure to heat is likely to have degraded the vaccine beyond an acceptable level.

The interpretation of the VVM is simple. Focus on the central square. Its color will change progressively. As long as the color of this square is lighter than the color of the ring, the vaccine can be used. As soon as the color of the central square is the same color as the ring or of a darker color than the ring, the vaccine should be discarded.

Shelf-Life

The suspension can be used within 24 months from the date of manufactured if kept as recommended.

Registration number

Euvichol-Plus Suspension Oral Cholera Vaccine– MAL19126001AZ

Manufactured by

Eubiologics Co., Ltd
Basement, 1F, 2-4~6 and 3-1~4 of 4-dong
56, soyanggang-ro, chuncheon-si
Gangwon-do, South Korea, 24232

Product Registration Holder

Averroes Pharmaceutical Sdn. Bhd.
03-08-01 & 03-09-01, Block 2, Presint Alami
Worldwide Business Center II
Persiaran Akuatik, Seksyen 13
40100 Shah Alam, Selangor
Malaysia
Tel : +603 5511 1433
Fax: +603 5511 1431

Date of revision of package insert

4th March 2025