

# Vimelinspec

Powder

## COMPOSITION:

Lincomycin (as Lincomycin hydrochloride).....44 g  
Spectinomycin  
(as Spectinomycin dihydrochloride).....44 g  
Exp. qs.....1000 g

## DESCRIPTION

The finished product is fine white powder

## PHARMACODYNAMICS

The lincosamide antibiotics lincomycin and clindamycin, share mechanisms of action and have similar spectrums of activity although lincomycin is usually less active against susceptible organisms. They may act as bacteriostatic or bactericidal agents, depending on the concentration of the drug at the infection site and the susceptibility of the organism. The lincosamides are believed to act by binding to the 50S ribosomal subunit of susceptible bacteria, thereby inhibiting peptide bond formation.

Most aerobic gram positive cocci are susceptible to the lincosamides (*Strep. faecalis* is not), including staphylococcus and streptococci. Other organisms that are generally susceptible include:

*Corynebacterium diphtheriae*, *Nocardia asteroides*, *Erysipelothrix*, and *Mycoplasma* spp. Anaerobic bacteria that may be susceptible to the lincomycin include: *Clostridium perfringens*, *C. tetani* (not *C. difficile*), *Bacteroides* (including many strains of *B. fragilis*), *Fusobacterium*, *Peptostreptococcus*, *Actinomyces* and *Peptococcus*.

Spectinomycin is primarily a bacteriostatic antibiotic that inhibits protein synthesis in susceptible bacteria by binding to the 30S ribosomal subunit.

Spectinomycin has activity against a wide variety of gram positive and gram negative bacteria, including *E. coli*, *Klebsiella*, *Proteus*, *Enterobacter*, *Salmonella*, *Streptococci*, *Staphylococcus* and *Mycoplasma*. It has minimal activity against anaerobes, most strains of *Pseudomonas*, *Chlamydia*, or *Treponema*.

## PHARMACOKINETICS

Oral absorption of lincomycin may be greatly reduced by the presence of food in the stomach. Oral absorption of pig is 20 to 50%.

Spectinomycin is only slightly absorbed from the gastrointestinal tract. 24 hours following oral administration, concentrations of spectinomycin are found in the following swine tissues in decreasing concentrations: kidney, liver, lung, muscle, and fat.

## INDICATION

Prevention and treatment of bacterial infections such as vibrio diarrhea, erythematosis, Salmonella, coliform enteritis, staphylococcus, streptococcosis, arthritis, swine monoclonal, mycoplasma pneumonia, haemophilus pneumonia, atrophic rhinitis, piglet diarrhea.

## TARGET SPECIES

Swine

## ROUTE OF ADMINISTRATION

To be mixed with feed.

## RECOMMENDED DOSE

Treatment: 1 ~ 3 kg of product per ton of feed for 1-3 weeks.

Prevention: 0.25 to 1kg of product per ton of feed.

## CONTRA-INDICATIONS

Do not use in case of hypersensitivity to lincomycin, spectinomycin.

Do not use on horses, rabbits, hamsters, guinea pigs and ruminating animals because of serious gastrointestinal effects that may occur, including death.

## INTERACTION WITH OTHER MEDICAL PRODUCT

Kaolin (found in several over-the-counter antidiarrheal preparations) has been shown to reduce the absorption of lincomycin by up to 90% if both are given concurrently. If both drugs are necessary, separate doses by at least 2 hours. Because lincomycin has been associated with severe diarrheas in several species, the need for an antidiarrheal product may indicate impending toxicity.

Lincomycin possesses intrinsic neuromuscular blocking activity and should be used cautiously with other neuromuscular blocking agents.

Antagonism has been reported when spectinomycin is used with chloramphenicol or tetracycline.

## WARNINGS AND PRECAUTIONS

### Warnings:

Do not use this product within 24 hours before slaughter.

### Usage Precautions:

- 1) Only administer the prescribed dosage.
- 2) Please observe the withdrawal period.
- 3) Observe the prescribed usage and dosage as it may cause weakness and economic loss due to misuse.
- 4) Do not administer the same drug while simultaneously administering this agent.
- 5) Do not use on animals that have had shock and hypersensitivity to this medicine.
- 6) In addition to applicable animals, gastrointestinal disorders may appear when used.
- 7) Occasionally, stool, skin redness and irritant behavior may occur after administration, but it will heal naturally within 5-8 days.

### Other precautions:

- 1) Do not use this product on humans as it is a veterinary drug.
- 2) Use only as directed by your veterinarian.
- 3) Do not use on animals other than the target species because the safety and effectiveness are unknown.
- 4) To prevent accidents, keep out of reach of children and keep away from animals.
- 5) Observe the prescribed storage method as it may cause a change in stability and effectiveness.
- 6) Do not use the product after the expiration date. Dispose of it safely.
- 7) Dispose of used containers and packaging safely and do not use them for other purposes.
- 8) In case of skin or eye contact, wash off with water thoroughly and consult a doctor immediately if any abnormality occurs.

## PREGNANCY AND LACTATION

Safe use during pregnancy has not been established, but neither has the drug been implicated in causing teratogenic effects.

## SIDE EFFECTS:

On rare occasions, some pigs may show anal swelling, diarrhea, irritable behavior, and

skin reddening are generally self-limiting within 5 to 8 days.

## SYMPTOMS AND TREATMENT OF OVERDOSE

No specific information was located on oral overdoses, but because the drug is negligibly absorbed after oral administration, significant toxicity is unlikely via this route.

## STORAGE CONDITION

Store in a cool dry place (<30°C). Strictly avoid light and heat exposure.

**SHELF LIFE:** 2 years

**SHELF LIFE AFTER OPENING:** 24 hours

**SHELF LIFE AFTER RECONSTITUTION:** 24hours

## WITHDRAWAL PERIOD

Do not use this product within 24 hours before slaughter.

**PACKAGING:** 1000 g

**REVISED DATE:** Oct 07, 2019