

PACKAGE INSERT FOR Deco 6% w/w Premix Powder

Brand / product name	Deco 6% w/w Premix Powder				
Name and strength of active substance	Composition per 1 kg: Decoquinat..... 60 g				
Product Description	White Powder				
Pharmacodynamics	Decoquinat is a 4-hydroxyquinoline antiprotozoal compound active against Eimeria spp. and Toxoplasma spp. Decoquinat inhibits the development of coccidia in the small intestine in the early part of the infective cycle, resulting in lower morbidity and mortality. The exact mode of action is unknown.				
Pharmacokinetics	The drug is administered by the oral route, the main site of action being within the gastro-intestinal tract. It is poorly absorbed by the target species and is largely eliminated in faeces unchanged. Consequently, tissue residues are low and deplete rapidly with time. Recovery of the administered material via excretion is essentially complete.				
Environmental Properties	N/A				
Indication	For the treatment and prevention of coccidiosis in lambs and calves. As an aid in the control of coccidiosis in lambs, by medication of ewe feed. As an aid in the prevention of abortions and perinatal losses due to toxoplasmosis by medication of ewe feed.				
Recommended Dosage	Species and indication	Method of administration	Recommended dose level	Recommended duration of treatment	
	Treatment of coccidiosis in lambs and calves and prevention of coccidiosis in lambs	In lamb and calf feed	1 mg decoquinat/kg bodyweight daily	At least 28 days	

Prevention of coccidiosis in calves and as an aid in prevention of coccidiosis in lambs	In calf and ewe feed	0.5 mg decoquinate/kg bodyweight daily	At least 28 days
As an aid in the prevention of abortions and perinatal losses due to toxoplasmosis	In ewe feed	2 mg decoquinate/kg bodyweight daily	Continuously for 14 weeks prior to lambing

Feed intake will depend on the clinical situation of the animal and on the season of the year. To ensure a correct dose, the concentration of active ingredient in the feed must be adjusted according daily feed intake. The following formula is recommended:

$$\text{Kg Deco 6\% w/w Premix Powder/T feed} = \frac{\text{Dose (mg active ingredient/kg body weight)} \times \text{Total weight of the animals to be treated (kg)}}{\text{Daily feed intake (kg)} \times 60}$$

1. Treatment of coccidiosis in lambs and calves and prevention of coccidiosis in lambs:

Feed continuously for 28 days when coccidiosis is expected to be a hazard. Medication may be continued if there is further identified risk beyond this period.

2. Prevention of coccidiosis in calves and as an aid in prevention of coccidiosis in lambs by medication of the ewe's feed:

Feed continuously for at least 28 days to ewes when oocyst shedding is likely to be a hazard to lambs (i.e. before, during or after lambing) or to calves when coccidiosis is likely to occur. The above provides good control of oocyst shedding from ewes under most conditions. In cases where a more severe challenge exists, double dosage should be used.

3. As an aid in the prevention of abortions and perinatal losses due to toxoplasmosis by medication of ewe feed:

Feed continuously for the last two-thirds of pregnancy (i.e. for the final 14 weeks prior to lambing).

On farms with a history of toxoplasmosis abortions, it may be economically beneficial to segregate susceptible ewes (e.g. bought in ewe-lambs) and administer medicated feed only to

	these animals, as the majority of older ewes will have been previously exposed to toxoplasma infection and will therefore be immune.
Mode of Administration	Mixed in feed
Contraindication	Do not use in cases of hypersensitivity to the active substance or to any of the excipients.
Warning and Precaution	(a) Special warnings for each target species The use of the product will maintain normal growth under conditions of coccidial challenge but does not improve growth of healthy lambs or calves. (b) Special precautions for use To help obtain even distribution in the final feed, it is recommended first to mix thoroughly at the rate of 1 part of the product to 3 parts of feed before blending into the final mix. In the preparation of pelleted feed, preconditioning temperatures of up to 80°C for short periods have been used and shown to have no effect on the product. A manufacturer authorised to incorporate at levels below 2 kg per tonne must be responsible for mixing when incorporation is less than 2 kg per tonne of final feed. i. Special precautions for use in animals. Medication of ewe rations may not prevent coccidiosis occurring in lambs and should only be given in conjunction with lamb medication. Do not mix with or into feeds containing any other anticoccidial. ii. Special precautions to be taken by the person administering the veterinary medicinal product to animals. When handling the product, prevent direct contact with the skin, avoid inhaling dust and wash hands after use. Do not eat, drink or smoke when handling the product. Only handle in a well ventilated area. iii. Other precautions This product is only authorised for use in medicated feedingstuffs or premixtures. In both cases it must be thoroughly mixed with feedingstuffs materials to ensure it is evenly distributed throughout the mixture.

	Any premixture containing this product must be thoroughly mixed with feedingstuffs materials to ensure that is evenly distributed throughout the final feed.
Interaction with other Medicaments	Do not mix with or into feeds containing any other anti-coccidial product.
Treatment on usage during pregnancy and lactation	<u>Pregnancy</u> Can be used during pregnancy. <u>Lactation</u> Can be used during lactation.
Maximum Residual Limit (MRL)	<u>Cattle and Sheep:</u> Muscle, liver, kidney, fat – 500 µg/kg
Adverse Effects/ undesirable Effects	None known. The frequency of adverse reactions is defined using the following convention: - very common (more than 1 in 10 animals treated displaying adverse reaction(s)) - common (more than 1 but less than 10 animals in 100 animals treated) - uncommon (more than 1 but less than 10 animals in 1,000 animals treated) - rare (more than 1 but less than 10 animals in 10,000 animals treated) - very rare (less than 1 animal in 10,000 animals treated, including isolated reports.
Overdose and treatment	Overdosage is unlikely with in-feed medication. Decoquate dosages of 4 mg/kg bodyweight in sheep and lambs and 6 mg/kg bodyweight in calves have been found to be well tolerated and without observable side effects.
Incompatibilities (for injection only)	N/A
Withdrawal period(s)	Cattle and sheep: Meat and offal: zero days. Milk: Not authorised for use in animals producing milk for human consumption.
Storage conditions	Store below 30°C (room temperature).
Dosage form and packaging available	Dosage form: Powder Packaging: 25 kgs

Name and address of contract manufacturer	N.S. Laboratory Sdn. Bhd. 54, Jalan S2C2, Green Technology Park, Seremban 2, 70300 Seremban, Negeri Sembilan Darul Khusus, Malaysia.
Name and address of registration holder	Weissen Company (M) Sdn. Bhd. Lot 105, No. 84, Jalan Velox 23, Taman Velox, 48000 Rawang, Selangor, Malaysia.
Date of revision of package insert	26.01.2026
Shelf life of veterinary products as packaged for sales	24 months
Shelf life of veterinary products as packaged for sales	24 hours.