

ENCLOR EAR DROPS 5% W/V

DESCRIPTION

Clear, mobile liquid, free from visible impurities. Colour: Colourless or with slight yellow tinge.

Active Substance

Each ml contains Chloramphenicol 50mg

PHARMACODYNAMICS

Chloramphenicol is a broad-spectrum antibiotic which has shown to have therapeutic activity against a wide range of gram-negative and gram-positive bacteria. Staphylococci and many other bacteria are known to develop a minimal resistance to chloramphenicol. In severe infection the topical use chloramphenicol should be supplemented with appropriate systemic therapy.

PHARMACOKINETICS

This preparation is a topical preparation meant for local therapeutic effect.

INDICATIONS

Enclor ear drops are indicated in the treatment of bacterial otitis extern by *Escherichia coli*, *Haemophilus Influenzae*, *staphylococci*, *Morax-Axenfeld diplobacillus*, *Proteus morgani* and *Pseudomonas pyocyanea*.

RECOMMENDED DOSAGE

Two or three drops are instilled into the ear three times daily. Treatment should be continued for a period as decided by the physician.

ROUTE OF ADMINISTRATION

Topical

CONTRAINDICATIONS

This medicine is contraindicated in patients with a history of hypersensitive reaction to chloramphenicol or any constituents of the otic drops.

WARNINGS & PRECAUTIONS

Prolonged use in some cases may result in overgrowth of non-susceptible organisms including fungi. Should superinfection occur, the drug should be discontinued and/or appropriate therapy instituted.

THIS PRODUCT MAY CAUSE BLOOD DYSCRASIAS.

INTERACTIONS WITH OTHER MEDICAMENTS

Not applicable.

PREGNANCY AND LACTATION

Chloramphenicol crosses the placenta but no problems have been reported with the use of ear drops. As with all drugs in pregnancy and during breast feeding, care should be taken in assessing the risk-benefits.

SIDE EFFECTS/ ADVERSE REACTIONS

May occasionally cause bone marrow hypoplasia, blood dyscrasia or hypersensitivity reactions.

SYMPTOMS AND TREATMENT FOR OVERDOSAGE

Overdose is unlikely to occur with this preparation. In case of excessive amount being instilled into the ear, it should be drained out by tilting the head to one side and wash the ear with warm water.

PACK SIZES

Plastic bottle of 5ml (10 bottles x 5ml)

STORAGE CONDITIONS

Keep container tightly closed in a dry place. Store at 2°C-8°C. Protect from light and freezing. Discard after 30 days from date of opening.

INSTRUCTIONS FOR USE

Apply 2 - 3 drops into the affected ear 3 times daily. Following administration of ear drops patients should be advised to lie down with the affected ear uppermost for a minimum of 10 minutes. After this time cotton wool may be inserted into the ear and normal activities resumed.

SHELF LIFE

Product should not be used beyond the expiry date imprinted on the product packaging.

REGISTRATION NUMBER: MAL19860033AZ

**CONTROLLED MEDICINE
UBAT TERKAWAL**

**KEEP MEDICINES OUT OF REACH OF CHILDREN
JAUHI UBAT-UBATAN DARI KANAK-KANAK**

For further information, please consult your doctor or your pharmacist.

Revision Number: 04

Revision Date: 01 Sep 2023

Lt-028.06 Product Registration Holder and Manufacturer
IDAMAN PHARMA MANUFACTURING SDN BHD (661901-P)
LOT 24 & 25, JALAN PERUSAHAAN LAPAN,
BAKAR ARANG INDUSTRIAL ESTATE, 08000 SUNGAI PETANI
KEDAH DARUL AMAN, MALAYSIA