

Further information can be obtained from your doctor or pharmacist.

ROYCE®

Product Holder/Manufactured by:
ROYCE PHARMA MFG. SDN. BHD. (650435-X)
PT 1663, Nilai Industrial Estate,
71800 Nilai, Negeri Sembilan, Malaysia.

Revision date: 250325

ROYCE®

CINNARIZINE TABLET 25MG (Cinnarizine - Antihistamine)

Presentation:

A white colour, round, shallow biconvex tablet, plain on one side and a break line on the other.

Content:

Each tablet contains Cinnarizine 25 mg.

Pharmacological Information:

Antihistamines diminish or abolish the main actions of histamine in the body, probably by occupying the receptor sites in the effector cells to the exclusion of histamine; they do not prevent the production of histamine.

The effects of histamine are considered to be mediated by 2 sets of receptors termed H1 and H2. Those effects which are mediated by the histamine H1 receptors include the contraction of smooth muscle and dilatation and increased permeability of the capillaries. The effects of histamines on vascular smooth muscle are mediated by the H2 as well as the H1 receptors. Other effects which are mediated by H2 include cardiac accelerating effects and in particular, the stimulating action of histamine on the secretion of gastric acid.

In general, antihistamines are readily absorbed from the gastrointestinal tract, metabolized in the liver, and excreted usually mainly as metabolites in the urine.

The prominent mode of inactivation of histamine in most animal species was by methylation by histamine methyltransferase. Antihistamines acting on H1 had been shown to stimulate or inhibit histamine methyltransferase in vitro according to concentration. Similar findings had been made for the H2 receptor antagonist burimamide and metiamide.

Many histamine H1 receptor antagonists have other pharmacological properties largely independent of antihistamine activity, including anticholinergic, adrenaline-enhancing and/or adrenal-in-antagonising, and serotonin-antagonising effects. Some of these properties are not generally of beneficial clinical significance and may be regarded as side effects. Most histamine H1 receptor antagonist cause depression of the central nervous system changing to stimulation with very large doses most have some local anesthetic properties.

The secretion of gastric acid is inhibited by a class of antihistamines, with a close structural relationship to histamine, termed histamine H2 receptor antagonists.

Histamine, when released from the tissues, causes allergic reactions. The numerous manifestations of such reactions include asthma, hay fever, vasomotor rhinitis, and angioneurotic oedema. Unless the agent precipitating the patient is carefully desensitised a cure cannot be expected. Adrenaline and other sympathomimetic agents, antihistamines, and corticosteroids all provide there as long as the cause of the cause of the allergy persists.

Indication:

Cinnarizine is used for the control of vestibular disorders such as vertigo, tinnitus, nausea, and vomiting as seen in Meniere's disease. Cinnarizine is also effective in the control of motion sickness.

Dosage and Administration:

Adult and children over 12 years:
Vestibular disorders: One tablet to be taken three times a day.
Motion sickness: One tablet to be taken 2 hours before travel and half tablet every 8 hours during journey.
Children 5 to 12 years: Half the adult dose.
Use in elderly - As for adults.

Contraindications:

None.

Precautions:

Cinnarizine may cause drowsiness; patients affected in this way should not drive or operate machinery. Avoid alcoholic drink.

Pregnancy and Lactation:

Pregnancy

The safety of Cinnarizine in human pregnancy has not been established although studies in animals have not demonstrated teratogenic effects. As with other drugs, it is not advisable to administer Cinnarizine in pregnancy.

Nursing mothers:

It is not known whether Cinnarizine is excreted in human milk; however, low concentrations of other anti-histamines have been detected in human milk. Because of the increased risk of antihistaminic side effects in infants, particularly in neonates, patient should not nurse while taking this drug.

Interaction With Other Medicaments:

Alcohol, tranquilizers, barbiturates, sedative-hypnotics, and other CNS depressants - Increases CNS depression.
MAO inhibitors - Increases anticholinergic effect.

Side Effects:

Rarely, allergic skin reactions have been reported which have responded to discontinuation of therapy.

Symptoms and Treatment of Overdose:

There is no specific antidote to Cinnarizine and, in the event of overdosage, gastric lavage is recommended.

Storage Condition:

Keep container tightly closed. Store in a dry place below 30°C. Protect from light.

Pack Size:

Plastic container of 30 tablets.
Blister pack of 1 x10's, 10 x10's, 50 x10's and 100 x10's.

Product Registration Number:

MAL19861200AZ