

AMOXICAP

VIAO29-var (MY)

DESCRIPTION

Amoxicap 250 Capsule:

Scarlet opaque and grey opaque capsule with "HD" printed on one end and "AX250" on the other end of the capsule.

Amoxicap 500 Capsule:

Scarlet opaque/yellow opaque capsule with "AX500" printed on one end and "HD" printed on the other end.

COMPOSITION

Amoxicap 250 Capsule:

Amoxycillin (as trihydrate) 250 mg/capsule

Amoxicap 500 Capsule:

Amoxycillin (as trihydrate) 500 mg/capsule

ACTIONS AND PHARMACOLOGY

Amoxycillin, a bactericidal antibiotic, inhibits bacterial cell wall synthesis by preventing cross-linkage of peptidoglycan chains which is necessary for bacterial cell wall strength and rigidity. Cell division and growth are inhibited and lysis of susceptible bacteria frequently occurs.

Oral absorption is not greatly influenced by the presence of food and excretion is mainly via renal and biliary systems.

INDICATIONS

For treatment of :

- Ear, nose and throat infections caused by *streptococci*, *pneumococci*, non-penicillinase - producing *staphylococci* and *H. influenzae*.
- Genitourinary tract infections caused by *E. coli*, *P.mirabilis*, *S. faecalis*.
- Skin and soft-tissues infections caused by *streptococci*, non-penicillinase - producing *staphylococci* and *E. coli*.
- Anogenital and urethral *gonorrhoea* caused by *N. gonorrhoea*.

CONTRAINDICATIONS

- Contraindicated in patients known to be sensitive to penicillin.
- Avoid in patients with infectious mononucleosis because of increased risk of skin rashes.

PRECAUTIONS

- As with any potent drug, periodic assessment of renal, hepatic and hematopoietic function should be made during prolonged therapy. The possibility of superinfections with mycotic or bacterial pathogens should be kept in mind during therapy. If superinfections occur (usually involving *Enterobacter*, *Pseudomonas* or *Candida*), the drug should be discontinued and/or appropriate therapy instituted.
- Caution in patients with known histories of allergy.
- Should be used with caution in pregnancy and lactation. Safety for use in pregnancy has not been established.
- Reduced doses may be required in patients with severe impaired renal function.

WARNINGS

Serious and occasionally fatal hypersensitivity (anaphylactoid) reactions have been reported in patients on penicillin therapy although anaphylaxis is more frequent following parenteral therapy, it has occurred in patients on oral penicillins. These reactions are more likely to occur in individuals with a history of sensitivity to multiple allergens.

There have been reports of individuals with a history of penicillin hypersensitivity who have experienced severe reactions when treated with Cephalosporins. Before therapy with any penicillin, careful inquiry should be made concerning previous hypersensitivity reactions to Penicillins, Cephalosporins or other allergens. If an allergic reaction occurs, appropriate therapy should be instituted and discontinuance of amoxicillin therapy considered. Serious anaphylactoid reactions require immediate emergency treatment with Epinephrine. Oxygen, intravenous steroids and

airway management, including intubation, should also be administered as indicated.

SIDE/ADVERSE EFFECTS

- Hypersensitivity reaction:
Rash and less frequently as exfoliative dermatitis or erythema multiforme.
- Hemolytic effects:
Hemolytic anemia / anemia / eosinophilia / leukopenia / neutropenia / agranulocytosis / thrombocytopenia / thrombocytopenic purpura.
- G.I. effects:
Diarrhoea, nausea, vomiting, black hairy tongue, glossitis, stomatitis, sore mouth or tongue.
- Renal effects:
Acute interstitial nephritis.
- Hepatic effects:
A moderate increase in serum concentration of AST (SGOT).
- Skin and subcutaneous tissue disorders:
Frequency 'very rare': Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS).

DRUG INTERACTIONS

- Concurrent use with Allopurinol or Probenecid requires careful monitoring.
- Concurrent use with Chloramphenicol, Erythromycins, Sulfonamides and Tetracycline may interfere with the bactericidal effect of amoxycillin.

OVERDOSE

Clinical features : Anorexia, nausea, vomiting, abdominal discomfort, diarrhoea.

Treat overdose by emesis or gastric lavage, if appropriate and other supportive measures.

DOSAGE AND ADMINISTRATION

Adults:

Oral, 250 to 500 mg every eight hours; or as directed.

Children:

Infants up to 6 kg of body weight:

Oral, 25 - 50 mg every 8 hours.

Infants 6 - 8 kg of body weight:

Oral, 50 - 100 mg every 8 hours.

Infants and children 8 - 20 kg of body weight:

Oral, 6.7 - 13.3 mg/kg of body weight every 8 hours.

Children 20 kg of body weight and over:

See adult dose.

Note: The information given here is limited. For further information consult your doctor or pharmacist.

Storage:

Amoxicap 250 Capsule

Store below 30°C. Protect from moisture.

Amoxicap 500 Capsule

Store below 30°C. Protect from moisture.

Presentation/Packing:

Capsule 250 mg x 100's, 500's, 1000's,

Blisters 10 x 10's, 100x10's

Capsule 500 mg x 100's, 500's, Blisters 10 x 10's

(Not all presentation is available locally)

Product Registration Holder / Manufactured by: HOVID Bhd.

121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar),

30010 Ipoh, Perak, Malaysia.

Revision date: November 2021

hovid