



PRINAZIN TABLET 25MG

Description: White, round, flat with top-scored tablet. Diameter = 7.00mm.

CONTENT

Cinnarizine 25mg/tablet

INDICATIONS

- Disorders of balance - maintenance therapy for symptoms of labyrinthine disorders, including vertigo, dizziness, tinnitus, nystagmus, nausea and vomiting.
- Prophylaxis of motion sickness.

PHARMACOLOGY

Cinnarizine has antihistamine activity and is also a smooth muscle relaxant and vasodilator. Sedative effects are not marked. It is well absorbed orally, metabolised in the liver, and excreted in the urine.

ADVERSE EFFECTS

Allergic skin reactions, fatigue, drowsiness and dyskinesia, extrapyramidal disorder, Parkinsonism and tremor in elderly people during prolonged therapy.

DRUG INTERACTIONS

Alcohol, CNS depressants and tricyclic antidepressants - The sedative effects of cinnarizine and of any of the following may be potentiated when used concomitantly.

Diagnostic interference - Because of its antihistamine effect, cinnarizine may prevent otherwise positive reactions to dermal reactivity indicators if used up to 4 days prior to skin testing.

PRECAUTIONS

- As with other antihistamines cinnarizine may cause epigastric distress; taking it after meals may diminish gastric irritation.
- In patients with Parkinson's disease cinnarizine should only be given if the advantages outweigh the possible risk of aggravating this disease.
- Cinnarizine may cause somnolence, especially at the start of treatment. Therefore, caution should be taken when alcohol, central nervous system (CNS) depressants or tricyclic antidepressants are used concomitantly.

PREGNANCY AND LACTATION

Safety for use in pregnancy and lactation has not been established.

Pregnancy: Although in animal studies, cinnarizine has shown no teratogenic effects, as with all drugs, cinnarizine should be used during pregnancy only if the therapeutic benefits justify the potential risks for the fetus.

Breast-feeding: There are no data on the excretion of cinnarizine in human breast milk: nursing should therefore be discouraged in women using cinnarizine.

DOSAGE AND ADMINISTRATION

Disorders of balance:

Adult: 1 tablet of 25 mg three times a day

Prophylaxis of motion sickness:

Adult and adolescent aged 13 years and above: 1 tablet of 25 mg at least half an hour before travelling; to be repeated every 6 hours;

Children aged 6 to 12 years: half of the adult dose is recommended.

Cinnarizine should preferably be taken after meals. It is for oral administration.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE

Caution when driving or doing jobs requiring alertness because of possible drowsiness.

TREATMENT OF OVERDOSAGE

Symptoms: The signs and symptoms are mainly due to the anticholinergic activity of cinnarizine. Acute overdoses have been reported with doses ranging from 90 to 2250 mg. The most common symptoms include alterations in consciousness ranging from somnolence to stupor and coma, vomiting, extrapyramidal symptoms, and hypotonia. In a small number of young children, seizures developed.

Clinical consequences were not severe in most cases, but deaths have been reported after single and polydrug overdoses involving cinnarizine. Treatment: Symptomatic and supportive care.

Storage

: Store in cool, dry place below 30°C. Protect from light. Keep out of reach of children.

Presentation/Packing

: Tablet 25mg x 100x10's ; 1000's(EXPORT)

Manufactured &

: Prime Pharmaceutical Sdn. Bhd.

Distributed by

1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia.