



ECOCIN CREAM

Description: White opaque cream

Contents

Econazole Nitrate	1% w/w
Triamcinolone Acetonide	0.1% w/w
Chlorocresol	0.1% w/w

Indications

For treatment of cutaneous candidiasis, pityriasis versicolor, and tinea accompanied by inflammatory or allergic symptoms.

Pharmacology

Econazole, an imidazole antifungal agent, damages cell membrane and alters its permeability resulting in loss of essential intracellular elements. Triamcinolone acetonide, a high-potency glucocorticoid acts by diffusing across cell membranes and stimulates transcription of mRNA resulting in protein synthesis of various enzymes ultimately responsible for its anti-inflammatory effects.

It is absorbed to a small extent percutaneously. Occlusion or use over extensive areas increases topical absorption.

Adverse Effects

Side/adverse effects may be more pronounced with the use of an occlusive dressing.

Local erythema, stinging or burning sensations may occur occasionally.

With prolonged use, moon face, thinning of skin with easy bruising, facial hair growth, loss of hair on scalp, reddish purple lines on arms, legs, trunk, or groin may occur.

Precautions

- Do not use in or around the eye.
- Do not bandage or wrap the treated skin area unless directed to do so by physician.
- Paediatric therapy continuing for longer than 2 weeks should be evaluated carefully by the physician.
- Caution in patients with tuberculosis of the skin and in nursing mothers.

Contraindications

- Not recommended for patients with systemic fungal infections, acute infections including herpes zoster and herpes simplex ulceration of the eyes and vaccination with live vaccine.
- Excessive or prolonged usage in pregnancy should be avoided.

Dosage and Administration

Apply sparingly to the affected area:

- | | |
|----------|---|
| Adults | — Twice a day, or as directed by physician. |
| Children | — Once a day, or as directed by physician. |

Overdosage

When applied to large areas, when the skin is broken, or under occlusive dressings, it may be absorbed in sufficient amounts to cause systemic effects.

Treatment consists of symptomatic measures, where possible the dosage being reduced or the drug slowly withdrawn.

Storage

Presentation/Packing

Manufactured & Distributed by

: Keep container tightly closed. Store in a cool, dry place, below 30C. Protect from light.
: Keep out of reach of children.
: 10gm / tube
: **PRIME PHARMACEUTICAL SDN. BHD.**
1505, Lrg Per. Utama 1, Tmn Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia.