



PRITALIN SYRUP 1.5MG / 5ML

Description: Plain and blackcurrant flavoured syrup.

Each 5ml contains : Terbutaline Sulphate 1.5mg
Methylparaben 0.1% w/v
Propylparaben 0.01% w/v

Indications:

For the treatment of asthma, chronic bronchitis, emphysema, and other bronchopulmonary disorders involving bronchospasm.

Pharmacology:

Terbutaline sulphate is a direct acting sympathomimetic amine with bronchodilator actions, thereby relieving bronchospasm and reducing airway resistance.

Basic parameters have been evaluated in man after i.v and oral administration of therapeutic doses e.g. IV single dose:

Volume distribution (vss): 114L

Total body clearance (cl): 213 ml/min

Mean residence time (mrt): 9.0 h

Renal clearance (clr): 149 ml/min (males)

Oral dose:

Renal clearance (clr): 1.925 ml/min (males)

Renal clearance (clr): 2.32 ml/min (females)

The plasma concentration/time curve after iv administration is characterised by a fast distribution phase, an intermediate elimination phase and a late elimination phase. Terminal half-life ($t_{1/2}$) has been determined after single and multiple dosing (mean values varied between 16 - 20 h). Bioavailability: Food reduces bioavailability following oral dosing (10% on average). Fasting values of 14 - 15% have been obtained. Metabolism: The main metabolite after oral dosing is the sulfate conjugate and also some glucuronide conjugate can be found in the urine.

Adverse effects / Side Effects:

Dizziness, drowsiness, muscle cramps or twitching, nausea, vomiting, unusual increase in sweating, palpitation, tachycardia, nervousness, headaches, restlessness, weakness and increase in blood pressure may occur in some patients.

Warning & Precaution:

Caution in patients with cardiovascular disorders, diabetes mellitus, hypertension, hyperthyroidism and history of seizures.

Check with physician immediately if difficulty in breathing persists after using this medication or if condition becomes worse.

Safety for use in pregnancy and lactation has not been established.

Tocolysis: Serious adverse reactions including death have been reported after administration of terbutaline / salbutamol to women in labor. In the mother, these include increased heart rate, transient hyperglycaemia, hypokalaemia, cardiac arrhythmias, pulmonary oedema and myocardial ischaemia. Increased fetal heart rate and neonatal hypoglycaemia may occur as a result of maternal administration.

Contraindications:

Not recommended for patients who are hypersensitive to other sympathomimetics.

Drug Interactions:

Co-administration of non-selective beta-adrenoceptor blocking agents may antagonise the bronchodilating effect of terbutaline.

Concurrent use with other sympathomimetics may increase the effects of either these medication or terbutaline and potential for side-effects.

Overdosage:

Symptoms: Headache, anxiety, tremor, nausea, tonic cramp, palpitations, tachycardia, arrhythmia, fall in blood pressure, hypokalaemia, hyperglycaemia and lactic acidosis.

Treatment: Reducing dosage in mild to moderate cases. In severe cases: gastric lavage, administration of activated charcoal, determination of acid-base balance, blood sugar and electrolytes, particularly serum potassium levels. Monitoring of the heart rate, rhythm and blood pressure. Metabolic changes should be corrected.

A cardioselective beta-blocker (e.g. metoprolol) is recommended for arrhythmias causing haemodynamic deterioration. The beta-blocker should be used with care due to the possibility of bronchoconstriction, caution in patients with a history of bronchospasm. If the beta2-mediated reduction in the peripheral vascular resistance significantly contributes to the fall in blood pressure, a volume expander should be given.

Dosage and administration:

Adults : Oral, 2 to 3 teaspoonfuls three times daily, administered at approximately 6 hour intervals, up to 10 teaspoonfuls per 24 hours.
Children 3 to 7years : Oral, ½ to 1 teaspoonful three times daily.
8 to 12 years : Oral, 1 to 2 teaspoonfuls three times daily.

Storage conditions : Keep container tightly closed. Keep out of reach of children.
Store in a cool, dry place below 30°C. Protect from light.

Presentation / Packing : Syrup 1.5mg / 5ml x 60ml ; 3.6L (EXPORT)

Manufactured & Distributed by : PRIME PHARMACEUTICAL SDN. BHD.
1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia.

28/04/2022