

LEAFLET

ASMALIN SYRUP 1.5MG/5ML

Each 5ml contains:
Terbutaline Sulphate 1.5 mg

Product Description:

A clear, colourless syrup with cherry flavour.

Pharmacodynamics:

Terbutaline Sulphate is a beta adrenergic receptor agonist which exert a preferential effect on beta₂ adrenergic receptors such as those located in bronchial smooth muscle. It has been shown in controlled clinical studies to relieve bronchospasm in chronic obstructive pulmonary disease.

Pharmacokinetics:

Terbutaline is incompletely absorbed from the gastro-intestinal tract and is also subject to fairly extensive first-pass metabolism by sulphate and some glucuronide conjugation in the liver and possibly the gut wall. Following the administration of terbutaline, a clinically significant improvement in pulmonary function occurs at 60 - 120 minutes. The maximum effect usually occurs within 120 - 180 minutes. Terbutaline also produces a significant decrease in pulmonary airway resistance which persist for at least four hours. Significant bronchodilation has been demonstrated for periods up to 8 hours. Terbutaline is excreted in the urine partly as the inactive conjugates and partly as unchanged terbutaline.

Indication:

Terbutaline Sulphate is indicated as a bronchodilator for bronchial asthma and for reversible bronchospasm which may occur in association with bronchitis or emphysema.

Recommended Dosage:

The usual oral dose for adults is 5mg administered at approximately six hourly interval, three times daily. A dose of 2.5mg three times daily is recommended for children of age between 12 - 15 years. In adults a total dose of 15mg should not be exceeded in a 24 hour period. In children a total dose of 7.5mg should be exceeded in a 24 hour period.

Route of Administration:

Oral

Contraindications:

Terbutaline Sulphate is contraindicated when there is known hypersensitivity to sympathomimetics.

Warnings and Precautions:

Tocolysis: Serious adverse reaction including death have been reported after administration of terbutaline/ salbutamol to women in labor. In the mother, these include increased heart rate, transient hyperglycaemia, hypokalaemia, cardiac arrhythmias, pulmonary oedema and myocardial ischaemia. Increased fetal heart rate and neonatal hypoglycaemia may occur as a result of maternal administration.

Terbutaline Sulphate should be used with caution in patient with diabetes, hypertension, hyperthyroidism and a history of seizures. As with other sympathomimetic bronchodilator agents, it should be administered cautiously to cardiac patients, especially those with associated arrhythmias. The concomitant use of terbutaline with other sympathomimetics is not recommended since their combined effect on the cardiovascular system may be deleterious to the patients.

Interactions with Other Medicaments:

β-blockers may inhibit bronchodilating effect.

Pregnancy and Lactation:

Although no teratogenic effects have been observed in animals or in patients, caution is recommended during the first trimester of pregnancy. Terbutaline Sulphate has been taken by a large number of pregnant women and women of childbearing age without any reported increase of malformation frequency or other signs of disturbance of the reproductive process. Terbutaline Sulphate passes over to maternal milk but the risk of influence on the child is unlikely with therapeutic doses.

Side Effects:

Commonly observed side effects included nervousness and tremor. Other reported reaction included headache, increased heart rate, palpitations, drowsiness, nausea, vomiting, sweating and muscle cramp. These effects are generally transient in nature.

Symptoms and Treatment of Overdose:

Excessive beta-adrenergic stimulation may augment the signs and symptoms listed under ADVERSE REACTIONS. Treat the alert patient by emptying the stomach by emesis, followed by gastric lavage. In the unconscious patient, secure the airway with a cuffed endotracheal tube before inducing lavage. Instillation of activated charcoal slurry may help reduce the absorption of Terbutaline Sulphate. Maintain adequate respiratory exchange. Provide cardiac and respiratory support as needed.

Effect on Ability to Drive and Use Machine:

Not known.

Storage Conditions:

Store at or below 30°C. Protect from light.

Pack Sizes:

A bottle of 60ml, 100ml and 120ml.

Pack Sizes (export only): A bottle of 3.6 litres and 3.8 litres.

Shelf-life:

3 years.

FURTHER INFORMATION CONCERNING THIS DRUG CAN BE OBTAINED FROM YOUR FAMILY PHYSICIAN / LOCAL GENERAL PRACTITIONER / PHARMACIST.

Manufacturer & Product Registration Holder:

Sunward Pharmaceutical Sdn. Bhd.

No. 9, 11 & 17, Jalan Kempas 4,
Taman Perindustrian Tampoi Indah,
81200 Johor Bahru, Johor, Malaysia.

SL303A-R9
Revision Date: 04/07/2025