

pharmaniaga®

TUSSINOL-N SYRUP

DESCRIPTION

Clear, mobile liquid, free from visible impurities. Colour: Green.

Each 5ml contains:

Brompheniramine Maleate 2mg
Phenylephrine Hydrochloride 5mg

Preservative: Sodium Benzoate 5mg

PHARMACODYNAMICS

Pharmacotherapeutic group: Adrenergic and dopaminergic agents

ATC code: R01BA03

Mechanism of action

Antihistaminic (H₁-receptor) : Antihistamines used in the treatment of allergy act by competing with histamine for H₁-receptor sites on effector cells. They thereby prevent, but do not reverse, responses mediated by histamine alone. The anticholinergic actions of most antihistamines provide a drying effect on the nasal and oral mucosa.

Decongestant: Sympathomimetic amines act on alpha-adrenergic receptors in the mucosa of the respiratory tract to produce vasoconstriction, which temporarily reduces the swelling associated with inflammation of the mucous membranes lining the nasal passages

PHARMACOKINETICS

Absorption : Brompheniramine is well absorbed after oral administration; Phenylephrine has reduced bio-availability (about 38%) from gastrointestinal tract because of first-pass metabolism by monoamine oxidase in the stomach and liver.

Metabolism : Antihistamine - Hepatic (cytochrome P-450 system) and some renal; Sympathomimetic amines - Phenylephrine undergoes extensive metabolism in the intestinal wall and in the liver. Sulfate conjugates are formed largely in the intestinal wall. Also, undergoes oxidative deamination by monoamine oxidase.

Half-life : Brompheniramine - 25 hours; Phenylephrine - 2.1 to 3.4 hours.

Onset of Action : Most first generation antihistamines - 15 to 60 minutes.

Time to peak concentration : Brompheniramine - 2 to 5 hours; Phenylephrine - 0.75 to 2 hours (to achieve peak level ranging from 0.9 to 298 ng/mL, respectively)

Time to peak effect : Brompheniramine - 3 to 9 hours

Duration of Action : 4 to 12 hours

Excretion : Antihistamine - Renal. Based on published data, most of antihistamines are excreted as metabolites within 24 hours.

Phenylephrine - 2.6% of the administered oral dose is excreted unchanged, 80 to 86% of unchanged Phenylephrine and metabolites are recovered in 48 hours after oral administration.

INDICATIONS

- Indicated for the relief of nasal congestion, hypersecretion and cough due to common cold.
- Symptomatic relief of allergic manifestations of upper respiratory tract disorders.
- Effectively controls post-nasal drip and its consequent cough.

RECOMMENDED DOSAGE

Adults : 2 teaspoonfuls (10mL) 4 to 6 hourly.

Children (2-6 years) : 1/2 teaspoonful (2.5mL) 4 to 6 hourly and should be used only as directed by a physician.

(6-12 years) : 1 teaspoonful (5.0mL) 4 to 6 hourly.

ROUTE OF ADMINISTRATION

Oral

CONTRAINDICATIONS

A history of hypersensitivity to the substance or substances of similar chemical structure (or any of the other ingredients in the product).

WARNINGS & PRECAUTIONS

- Patients with renal or hepatic impairment and patients with epilepsy should use with caution.
- Phenylephrine has an excessive vasopressor response that may cause a prolonged rise in blood pressure. It induces tachycardia or reflex bradycardia and should be avoided in severe hyperthyroidism. Patient with severe ischaemic heart disease should use with caution.
- Phenylephrine should be avoided in patients with diabetes mellitus or prostatic hyperplasia.
- Not recommended for use in newborn or premature infant. Older children may have a paradoxical reaction characterized by hyperexcitability.
- When used for treatment of cough and cold:
 - Not to be used in children less than 2 years of age
 - To be used with caution and doctor's/pharmacist's advice in children 2 to 6 years of age.

INTERACTIONS WITH OTHER MEDICAMENTS

The following interactions have been noted: Central Nervous System (CNS) depressants (alcohol, sedatives, opioid analgesics, hypnotics);

- May cause an increase in sedation effects monoamine oxidase inhibitors (MAOIs) and tricyclic antidepressants (TCAs);
- May prolong and intensify the anticholinergic and CNS depressive effects other sympathomimetic agents, such as decongestants, appetite suppressants and amphetamine-like psychostimulants;
- May cause an increase in blood pressure and additive effects methyl dopa and Beta blockers;
- May cause an increase in blood pressure.

PREGNANCY & LACTATION

- Tussinol-N Syrup should be used in pregnancy only if the potential benefits to the patient outweighs the possible risk to the fetus.
- Small amounts of antihistamines are excreted in breast milk and its use is not recommended in nursing mothers unless the potential benefits to the patient outweighs the possible risk to the infant.

SIDE EFFECTS

Central Nervous System (CNS)

CNS depressive effects include sedation and impaired

performance (impaired driving performance, poor work performance, reduced motor skills and impaired information processing). Performance may be impaired in the absence of sedation and may persist the morning after a night-time dose. CNS stimulatory effects may include anxiety, hallucinations, appetite stimulation, muscle dyskinesia and activation of epileptogenic foci.

Anticholinergic effects

Adverse effects related to cholinergic blockage include dryness of the eyes and nose, blurred vision, urinary retention and hesitancy, constipation and tachycardia.

Others

Skin rashes and urinary retention in elderly and children are more likely to experience side effects.

SYMPTOMS & TREATMENT FOR OVERDOSAGE

Symptoms:

Nervousness, tremor, insomnia, agitation, irritability, abdominal pain, severe headache, nausea and vomiting.

Treatment:

Stomach should be emptied by aspiration and lavage. Diazepam can be given to control CNS stimulation and convulsions. Alpha-adrenergic blockers are useful to manage hypertension.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINES

This product may cause drowsiness. Drowsiness may continue the following day. Alcohol should be avoided as it may increase drowsiness. Those affected should not drive or operate machinery.

PACK SIZES

Plastic Containers of 120mL.

STORAGE CONDITIONS

Keep container tightly closed in a dry place, below 30°C; Protect from light.

INSTRUCTIONS FOR USE

Take with or without food.

Do not take with alcoholic drinks.

SHELF LIFE

Product should not be used beyond the expiry date imprinted on the product packaging.

REGISTRATION NUMBER

MAL19910130AZ

CONTROLLED MEDICINE UBAT TERKAWAL

KEEP MEDICINE OUT OF REACH OF CHILDREN JAUHI UBAT-UBATAN DARI KANAK-KANAK

For further information, please consult your doctor or your pharmacist.

Revision Number : 05
Revision Date : 25 Nov 2025

LT-052.06 Product Registration Holder and Manufacturer

INSPIRAPHARMA SDN. BHD. (200401023395)
LOT 24 & 25, JALAN 8,
KAWASAN PERUSAHAAN BAKAR ARANG,
08000 SUNGAI PETANI,
KEDAH DARUL AMAN, MALAYSIA.