

PROZYME TABLET (Enzyme)

Presentation:

A round, flat face, bevel edged green coloured tablet, with a breaking line on one side, 7.9mm in diameter and with peppermint flavour.

Content:

Each tablet contains:

Papain 2.25 W.P.A unit

(equivalent to 150,000 USP unit).

Pharmacological Information:

Papain is a proteolytic enzyme which hydrolyses polypeptides, amides and esters, especially at bonds involving basic amino acids or leucine or glycine, yielding peptides of lower molecular weight. Thus reduces the swelling at the inflammatory site.

Papain is an activated and purified preparation of protease extracted from the topical fruit Carica Papaya. The optimum pH of papain is 5.7 at 37°C with a range 3.0-9.0 at which the activity is still 50%. One unit of Wallerstein Papain Activity (W.P.A) is defined as the quality of enzyme that will digest 1.25 g Hammarsten casein up to 55% within an hour at 40°C in the test condition. The term W.P.A expresses the number of proteasic activity units contained in 1 gram of a given enzymatic preparation.

Indication:

Acts as an antiinflammatory enzyme for use in conjunction with physical and/or chemotherapeutic measures for treatment or prevention of oedema and inflammatory conditions.

Dosage and Administration:

For oral use only.

2 tablets to be sucked or chewed 4 times daily for the first day, then 1 tablet 4 times daily for at least 5 days.

Contraindication:

Contraindicated in patients with known hypersensitivity to papain.

Warning and Precautions:

Concomitant use with anticoagulants, in patients with disorders of blood coagulation. Papain is very liable to deterioration. Its proteolytic activity is considerably impaired by oxidation.

Pregnancy and Lactation:

Safe use of this drug in pregnancy has not been established.

Side Effects:

Allergic reactions have followed repeated inhalations of papain powder. Urticaria, pruritus, dizziness, nausea, vomiting and diarrhoea have been observed following papain administration and it is possible that sensitisation may occur.

Symptoms and Treatment of Overdose:

None known.

Storage Condition:

Keep container tightly closed. Store in a dry place (below 30°C). Protect from light.

Pack Size:

Plastic container containing 50 tablets.

Blister pack of 1x10's, 10x10's, 50x10's and 100x10's.

Product Registration Number:

MAL19910323X

Further information can be obtained from your doctor or pharmacist.

The logo for ROYCE, featuring the word "ROYCE" in a bold, white, serif font with a registered trademark symbol (®) to its upper right, set against a solid black rectangular background.**Product Holder/Manufactured by:**

ROYCE PHARMA MFG. SDN. BHD. (650435-X)

PT 1663, Nilai Industrial Estate,

71800 Nilai, Negeri Sembilan, Malaysia.

Revision date: 280618