

DENZO TABLET 5MG

MAL19910626X

DESCRIPTION:

TABLET

Colour : Buff (Light Mandarin)
Shape : Round and Biconvex
Coating : Enteric-coated

EACH TABLET CONTAINS:

Serratiopeptidase 5 mg

PHARMACODYNAMICS:

- Potent bradykinin-decomposing activity
Bradykinin, phlogogenic polypeptide, is locally formed in the presence of inflammation, causing inflammatory symptoms and dilatation and enhancing permeability of capillary blood vessels, migration of leukocytes and pain. Serratiopeptidase has been shown to exert a bradykinin decomposing effect, which is interpreted as representing a mechanism of anti-inflammatory activity.
- Fibrinolytic activity
As one of the anti-inflammatory mechanism of anti-inflammatory enzymes, the enzymes themselves are considered to relieve the inflammatory symptoms by breaking down the protein or fibrin which clog the vascular wall of inflammatory focus and the tissues and then by normalising the absorption of inflammatory exudate.
- Enhancement of Antibiotic action
Accumulation of fibrin and presence of necrotic tissues, disturb transport of the antibiotic to the locally of the morbid focus, leading to less therapeutic effect. Serratiopeptidase helps to absorb or remove the necrotic tissues and breakdown products, promotes transport of the antibiotics to the lesion and thereby enhance the action of the latter.

PHARMACOKINETICS:

It is absorbed after oral administration and distributes throughout the body.

INDICATION:

As concomitant therapy for inflammatory oedema and swelling.

RECOMMENDED DOSE:

Adults: 2 tablets. Children: 1 tablet.
To be taken 3 times a day.

INDIKASI:

Digunakan untuk melegakan keadaan inflamasi dan bengkak.

CADANGAN DOS:

Dewasa: 2 biji. Kanak-kanak: 1 biji.
Diambil 3 kali sehari.

功能：解除病發症引起之各种炎腫。

服法：成人，2片。兒童，1片。每日3次。

JAUHI UBAT DARIPADA KANAK-KANAK
KEEP OUT OF REACH OF CHILDREN

遠離兒童

ROUTE OF ADMINISTRATION:

FOR ORAL USE ONLY

CONTRAINDICATIONS:

It is contraindicated in patients hypersensitive to the ingredients of the preparations.

WARNING AND PRECAUTIONS:

In patients with blood coagulation abnormalities severe hepatic and renal disturbances or under treatment with anti-coagulants, DENZO should be given under careful observation.
DENZO may occasionally cause gastrointestinal disturbances such as anorexia, gastric discomfort and nausea. Administration should be discontinued when hypersensitive symptoms, such as eruption, appear.

SIDE EFFECTS:

With the recommended directions, side-effects are unlikely to occur.

PACKING/PACK SIZE(S):

Blister-pack of 2x10's, 10x10's and 100x10's.
Blister pack of 10's/Blister Envelope and 20 Blister Envelopes/Box.

MANUFACTURER/PRODUCT REGISTRATION HOLDER:

DYNAPHARM (M) SDN. BHD. 198001011897 (65683-V)
2497, Mk. 1, Lorong Perusahaan Baru 5,
Kawasan Perusahaan Perai 3,
13600 Perai, Pulau Pinang, Malaysia.

PI3NT D001-05
Latest Revision: Nov 2022