

DYNA KAOLIN-PECTIN

MAL19910640X

DESCRIPTION:

SUSPENSION

Colour : Pink
Flavour : Cherry

CONTENT:

Each 30 ml contains:

Kaolin Light 6 g
Pectin 150 mg

Preservatives: Methyl Paraben 0.1% w/v, Propyl Paraben 0.01% w/v & Sodium Benzoate 0.1% w/v

PHARMACODYNAMICS:

Kaolin Light is an adsorbent. It is administered by mouth in the symptomatic treatment of enteritis, colitis and dysentery associated with food poisoning and alkaloidal poisoning. It may be administered suspended in water. The Kaolin adsorbs irritants and forms a protective coating on intestinal mucosa.

Pectin acts to consolidate the stool. It is an efficient emulsifying and gelling agent in acid media and may be used in the treatment of diarrhoea.

PHARMACOKINETICS:

The preparation combines the adsorbent and detoxifying effects of Kaolin (which functions principally in the stomach and small intestine) and Pectin (which exerts mainly in the large intestine).

INDICATION:

For the relief of minor diarrhoea.

RECOMMENDED DOSE:

Adults: 2 tablespoonfuls.

Children 7 to 12 years: 1 tablespoonful.

All to be taken 3 to 4 times daily.

CONTRAINDICATION:

The preparation should not be given to patients with intestinal obstruction, and it may be contraindicated in patients with spastic bowel conditions.

WARNING AND PRECAUTIONS:

Appropriate fluid and electrolyte therapy should be given to protect against dehydration. Oral rehydration therapy (which is the use of appropriate fluids including oral rehydration salts) remains the most effective treatment for dehydration due to diarrhoea. The intake of as much of these fluids as possible is therefore imperative.

KAOLIN/PECTIN/KAOLIN-PECTIN IS NOT RECOMMENDED FOR CHILDREN UNDER 6 YEARS OF AGE.
--

Severe constipation, which may lead to faecal impaction, may rarely occur in children and the elderly patients taking Kaolin and Pectin.

Kaolin and Pectin may interfere with the absorption of other drugs, including antibiotics, administered concurrently.

DRUG INTERACTION:

The adsorbent properties of Kaolin and other clays may influence the gastrointestinal absorption of other drugs. The absorption of Lincomycin has been reported to be significantly reduced by kaolin.

PREGNANCY AND LACTATION:

The safety of Kaolin during pregnancy and lactation has not been established, though no special precautions are thought necessary provided the recommended dose is taken.

SIDE EFFECTS/ADVERSE REACTIONS:

Large quantities of Pectin and other bulk laxative may temporarily increase flatulence and distension and intestinal obstruction has been reported after their administration. Oesophageal obstruction has also occurred when these agents have been swallowed dry.

PACKING/PACK SIZE(S):

Plastic bottle of 120 ml.

**JAUHI UBAT DARIPADA KANAK-KANAK
KEEP OUT OF REACH OF CHILDREN
SHAKE WELL BEFORE USE**

MANUFACTURER/PRODUCT REGISTRATION HOLDER:

DYNAPHARM (M) SDN. BHD. 198001011897 (65683-V)

2497, Mk. 1, Lorong Perusahaan Baru 5,
Kawasan Perusahaan Perai 3,
13600 Perai, Pulau Pinang, Malaysia.

PI2NL D003-01
Latest Revision: Dec 2021