



PATIENT INFORMATION LEAFLET

PLEASE READ THIS LEAFLET CAREFULLY BEFORE YOU TAKE THIS MEDICINE. IF YOU ARE UNSURE ABOUT ANYTHING ASK YOUR PHARMACIST OR DOCTOR.

WHAT IS FYBOGEL ORANGE?

Each sachet of Fybogel Orange holds a single dose of granules containing 3.5g ispaghula husk BP as the active ingredient. Each sachet also contains citric acid, potassium bicarbonate, sodium bicarbonate, polysorbate 80, silica colloidal anhydrous and orange flavour. Fybogel Orange is coloured with beta carotene and riboflavine sodium phosphate and is sweetened with aspartame and saccharin sodium. The granules are mixed with water to form an orange-flavoured drink.

Fybogel Orange comes in packs containing either 10 or 30 sachets.

The ispaghula husk in Fybogel Orange is a natural plant material which is high in fibre. It works by absorbing water and increasing the bulk of the food passing through the digestive system, helping your body to work normally.

WHY SHOULD I TAKE FYBOGEL ORANGE?

A lack of fibre in the diet is one of the main causes of bowel irregularity. Fybogel Orange is high in natural fibre which will help you to restore and maintain your body's regularity.

Fybogel Orange is also recommended wherever increased fibre is needed in your diet to help relieve constipation - this includes constipation during pregnancy or when breastfeeding. It is also recommended to help maintain a normal bowel function among patients suffering from any of the following conditions:-

- Irritable bowel syndrome
- Colostomy or ileostomy
- Haemorrhoids (piles)
- Anal fissure
- Persistent diarrhoea associated with diverticular disease
- Ulcerative colitis

BEFORE TAKING FYBOGEL ORANGE

As with all medicines, Fybogel Orange may not be suitable for some people.

Do not take Fybogel Orange if:

- You are allergic to Ispaghula husk or any of the other ingredients in this product
- You have a sudden change in bowel habit that has persisted for more than two weeks.
- The speed in which material moves through your gut has always been slow (reduced gut motility)
- your doctor has told you that you are suffering from a blockage of the bowel (intestinal obstruction or faecal impaction) or muscle weakness of the bowel wall (colonic atony)
- you have difficulty in swallowing or any throat problems
- you have phenylketonuria, because the product is sweetened with aspartame.
- You are suffering from an undiagnosed rectal bleeding and unable to pass stools following the use of laxatives.

If symptoms persist, consult your doctor.

Take special care:

- If you are a healthcare worker or care giver as allergic sensitisation may occur due to inhalation
- If not taken with enough liquid, can cause chest pain, vomiting or difficulty swallowing
- Elderly and debilitated (weak) patients may require medical supervision

Other medicines and Fybogel Orange:

- You must not take this medicine if you are taking other medicines which may cause constipation (e.g. codeine or morphine).
- if you are diabetic, you should take this product with medical supervision because adjustment of anti-diabetic therapy may be necessary.
- medicines for thyroid problems - your doctor may need to change the dose of the thyroid hormones you are taking.

HOW SHOULD I USE FYBOGEL ORANGE?

The amount of Fybogel Orange which should be taken depends upon age:

- Adults, including the elderly, and children over 12 years old, should take one sachet of Fybogel Orange twice a day, one in the morning and a second in the evening.
- Children aged between 6 and 12 years old should take half to one level 5 ml spoonful, depending on age and size, of Fybogel Orange twice a day, one in the morning and a second in the evening.
- Do not give Fybogel Orange to children under 6 years old.

Stir the required amount of Fybogel Orange into a glass of cold water (approximately 1/4 pint or 150 ml) and drink straight away. Fybogel Orange is best taken after meals.



Fybogel[®] ORANGE Ispaghula Husk



If the constipation does not resolve within 3 days or if abdominal pain occurs or in case of any irregularity of faeces, the use of ispaghula husk should be discontinued and medical advice must be sought.

Please consume a large amount of fluid/ water when taking this product.

Do not take Fybogel Orange without water. The product should be taken during the day at least 30 mins to 1 hour before or after intake of other medicines and should not be taken immediately before going to sleep. The effects start 12-24 hours later.

If you accidentally take too much, drink plenty of water.

Unless your doctor has given you different instructions on how much Fybogel Orange to take, follow the advice given in this leaflet.

WHAT SIDE-EFFECTS MAY OCCUR?

Like all medicines, this medicine can cause side effects, although not everybody gets them. This product can cause allergic reactions with symptoms including difficulty in breathing, wheezing or coughing, skin rash, itching and a sudden life-threatening allergic reaction. This product contains allergens and it is possible to come into contact with them via skin, inhalation or when taking the product by mouth and may cause red or itchy eyes, runny or blocked nose and a skin rash. If symptoms appear, you should immediately contact your doctor or pharmacist.

Due to the increased fibre intake of your diet you may experience flatulence or bloating during the first few days of taking Fybogel Orange. These effects should lessen as you continue to take Fybogel Orange.

If you have any other symptoms after taking this product, tell your doctor or pharmacist.

STORAGE

Do not use after the expiry date shown on the pack (EXP month/year). Keep Fybogel Orange in a dry place below 30 C.

Manufacturer and MA Holder: Reckitt Benckiser Healthcare (UK) Ltd., Hull, HU8 7DS, United Kingdom.

Fybogel Orange and  are trademarks.
Leaflet last revised May 2022





ข้อมูลนี้เป็นข้อมูลโดยย่อ
หากมีข้อสงสัยให้ปรึกษาแพทย์หรือเภสัชกร