

1. NAME OF THE MEDICINAL PRODUCT

ARCALION 200 mg, coated tablet

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Sulbutiamine 200mg for one coated tablet

Excipients with known effect: glucose, lactose, sunset yellow FCF aluminium lake (E 110), sucrose

For a full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Coated tablet.

4. CLINICAL PARTICULARS

4.1. Therapeutic indications

Symptomatic treatment of all forms of asthenia

4.2. Posology and method of administration

Posology

For adult use only: 2 to 3 tablets a day.

Tablets should be swallowed whole with a large glass of water, dividing the doses between the morning and midday meals.

Duration of treatment is limited to 4 weeks.

Paediatric population

The safety and efficacy of ARCALION have not been established in children and adolescents, ARCALION should not be used in this population.

Method of administration

Oral use.

4.3. Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

4.4. Special warnings and precautions for use

This medicinal product contains lactose, glucose and sucrose. This medicine should not be used by patients with galactose or fructose intolerance, Lapp lactase deficiency or glucose and galactose malabsorption syndrome (rare hereditary diseases) or a sucrase/isomaltase deficiency.

This medicine contains an azo colouring agent (E110) and may provoke allergic reactions (see section 4.8).

4.5. Interaction with other medicinal products and other forms of interaction

No specific interaction studies have been performed with ARCALION.

Concomitant use to be taken into consideration:

- **Diuretics:**

The urinary excretion of thiamine (metabolite of sulbutiamine) is increased.

- **Neuromuscular blocking agents:**

The effect of these medicines may be increased when administered concomitantly with thiamine (metabolite of sulbutiamine).

4.6. Fertility, pregnancy and breastfeeding

Pregnancy

There is no or limited data available (less than 300 pregnancy outcomes) on the use of sulbutiamine in pregnant women.

Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity (see section 5.3). As a precautionary measure, it is preferable to avoid the use of ARCALION during pregnancy.

Breastfeeding

It is unknown whether sulbutiamine/metabolites are excreted in human milk.

A risk to the newborns/infants cannot be excluded.

ARCALION should not be used during breast-feeding.

Fertility

There is no available data regarding the effects of sulbutiamine on fertility.

4.7. Effects on ability to drive and use machines

No specific studies on the effect on ability to drive and to use machines have been performed.

4.8. Undesirable effects

Tabulated list of undesirable effects

The following undesirable effects or events have been reported and are ranked using the following frequency: very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$); not known (cannot be estimated from the available data).

MedDRA system organ class	Frequency	Undesirable effects
Psychiatric disorders	Uncommon	Agitation
Nervous system disorders	Uncommon	Headache
		Tremor
Gastrointestinal disorders	Uncommon	Nausea
		Vomiting
	Not known	Gastralgia
		Diarrhoea
Skin and subcutaneous tissue disorders	Uncommon	Rash
General disorders and administration site conditions	Uncommon	Malaise

Description of selected undesirable effects

Due to the presence of sunset yellow FCF (E110), risk of allergic reactions (see section 4.4).

4.9. Overdose

Symptoms

In case of overdose: agitation with euphoria and tremor of the extremities. These symptoms are transient.

Treatment

Overdose symptoms can be treated according to the doctor's assessment.

5. PHARMACOLOGICAL PROPERTIES

5.1. Pharmacodynamic properties

Pharmacotherapeutic group: Vitamin B1, plain, ATC code: A11DA02

In humans:

This medicine has been studied in controlled clinical trials *versus* placebo or reference products, by using psychometric evaluation scales and tests. These trials demonstrate the psychoactive effect of this drug, with a predominant action on psychological and physical inhibition.

5.2. Pharmacokinetic properties

Sulbutiamine is rapidly absorbed and peak blood concentration is reached between 1 and 2 hours after administration.

The half-life is approximately 5 hours. Elimination is by the urinary route.

5.3. Preclinical safety data

No special hazards have been identified for use in humans on the basis of preclinical studies of acute, subchronic and chronic toxicity and reproduction toxicity (studies conducted in pregnant mice, rats and rabbits haven't shown any teratogenic potential). Sulbutiamine was not mutagenic in the Ames test. No studies of carcinogenicity have been performed.

6. PHARMACEUTICAL PARTICULARS

6.1. List of excipients

Maize starch, starch pregelatinised, anhydrous glucose, lactose monohydrate, magnesium stearate, talc, sodium hydrogen carbonate, carmellose sodium, white beeswax, titanium dioxide (E 171), ethylcellulose, sunset yellow FCF aluminium lake (E 110), glycerol oleate, polysorbate 80, povidone, sucrose, colloidal anhydrous silica.

6.2. Incompatibilities

Not applicable.

6.3. Shelf life

2 years

6.4. Special precautions for storage

Store below 30 °C.

6.5. Nature and contents of container

Aluminium canister of 30 tablets.

6.6. Special precautions for disposal and other handling

No special requirements.

7. MANUFACTURER :

Les Laboratoires Servier Industrie

45520 Gidy - France

8. PRODUCT REGISTRATION HOLDER:

Servier Malaysia Sdn Bhd

Unit No. 25-02, Level 25, Imazium

No.8, Jalan SS21/37, Damansara Uptown

47400 Petaling Jaya, Selangor Darul Ehsan

9. DATE OF REVISION

14 Jun 2022