

Package insert

hovid

hovid-Liquid Paraffin

VILIQ01-var_3

DESCRIPTION

Transparent, colourless, oily liquid, free from fluorescence by daylight, odourless or almost odourless and giving not more than a faint odour when heated.

COMPOSITION

Liquid Paraffin

PHARMACODYNAMICS

hovid-Liquid Paraffin, when taken internally, acts as a lubricant and keeps the stools soft. It softens fecal contents by coating them and thus preventing colonic absorption of fecal water. The lubricant effect eases passage of contents through intestines.

PHARMACOKINETICS

Absorption

It may be absorbed to a slight extent from the gastrointestinal tract, especially if emulsified.

INDICATIONS

hovid-Liquid Paraffin is indicated for the symptomatic relief of constipation. Long-term use of this medication is not recommended.

CONTRAINDICATIONS

- Laxatives should not be given to patients with intestinal obstruction, abdominal pain, nausea or vomiting, faecal impaction, colostomy and ileostomy.
- Mineral oil is not recommended for administration to children up to 3 years of age or to elderly patients because of the greater possibility of aspiration.

WARNING AND PRECAUTIONS

- Long-term use of this medication is not recommended.
- Safety for use in pregnancy and lactation has not been established. Not recommended for use in pregnant women.
- Repeated use is not advised.
- Consult your doctor if laxatives are needed every day, if you have persistent abdominal pain or a condition which makes swallowing difficult.

PREGNANCY AND LACTATION

None known

INTERACTION WITH OTHER MEDICAMENTS AND OTHER FORMS OF INTERACTION

Prolonged ingestion may interfere with the absorption of fat-soluble vitamins, and should be avoided.

MAIN SIDE/ADVERSE EFFECTS

Aspiration, with subsequent pulmonary infiltration has been reported.

OVERDOSE

Symptoms:

Anal seepage and irritation.

Treatment:

Discontinue therapy and consult physician immediately if overdose occurs.

DOSAGE AND ADMINISTRATION

Adults:

Orally, 15 to 45ml once a day, the dose being adjusted as needed.

Children 6-12 years:

Orally, 10 to 15ml once a day, the dose being adjusted as needed.

Note : The information given here is limited. For further information, consult your doctor or pharmacist.

Storage : Store below 30° C.

Protect from light.

Presentation/Packing : Bottle of 100ml, 3.6L

Product Registration Holder/Manufactured by : Hovid Bhd.,
121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar),
30010 Ipoh, Perak, Malaysia.

Revision date : December 2018.

hovid