

WINTOSAL CREAM

VIWIN03-var (MY)

DESCRIPTION

Pale green, opaque cream.

COMPOSITION

Methyl salicylate	15.0% w/w
Menthol	2.3% w/w
Camphor	4.0% w/w

ACTIONS AND PHARMACOLOGY

Methyl Salicylate is used externally as a counterirritant-rubefacient. When it is rubbed on the skin, it produces mild irritation and is applied in liniments and creams for the relief of pain in lumbago, sciatica and rheumatic conditions.

Camphor is used for its local action as a rubefacient when rubbed on the skin. When it is not applied vigorously, however, it may produce only a feeling of coldness. Camphor also has a mild local anaesthetic actions, and its application to the skin may be followed by numbness. Applied externally, camphor is used as a counterirritant in the treatment of sprains, fibrositis and neuralgia.

Applied to the skin menthol dilates the vessels, causing a sensation of coldness followed by an analgesic effect. It has been used for rheumatic pains and neuralgia when applied as liniments or creams. Counterirritants are drugs that act locally on cutaneous or mucosal tissue to produce "inflammation". The first response to local irritation is an increased circulation to the injured part. The localized vasodilation, mediated by way of an axon reflex, is attended by the feeling of comfort, warmth, and sometimes itching. Localized hyperesthesia also occurs. Drugs that evoke only reactive hyperemia are known as rubefacients.

PHARMACOKINETICS

Methyl salicylate readily absorbs through the skin.

Note: There are no data on the other ingredients in humans.

INDICATIONS

For temporary relief of pain associated with musculoskeletal soreness and discomfort.

WARNINGS

- This product is contraindicated in **children** under 2 years of age. Caution must be exercised when older children are treated.
- This product contains methyl salicylate and when applied or rub on to the skin, can be absorbed through the skin into the blood. For patients taking warfarin, excessive application to the skin for muscle or joint pains may increase the chances of bleeding.
- If you are taking warfarin, please consult your doctor before using this product.
- Do not use during pregnancy or when breastfeeding.

PRECAUTIONS

- It is dangerous to place any camphor-containing product into the nostrils of children. A small amount applied this way may cause immediate collapse.
- Avoid contact with the eyes.
- Do not apply to wounds or damaged skin.

MAIN SIDE/ADVERSE EFFECTS

Hypersensitivity reactions, urticaria and angioneurotic oedema may occur in susceptible persons.

OVERDOSE AND TREATMENT

Clinical features:

Hyperventilation, hyperthermia, epigastric pain, nausea, vomiting, vertigo, muscular excitability, confusion, respiratory alkalosis, metabolic acidosis, pulmonary oedema, hepatic and renal failure, convulsions and coma.

Treat poisoning by emesis or gastric lavage. Electrolyte and metabolic abnormalities, if any, should be corrected. Convulsions may be controlled by IV diazepam 5 - 10 mg.

DOSAGE AND ADMINISTRATION

Gently massage on intact skin two to three times a day. Best used after a hot bath.

Note: The information given here is limited. For further information, consult your doctor or pharmacist.

Presentation/Packing:

Cream x 15g and 30g
(Storage: Store below 30°C. Protect from light.)

Cream x 500g
(Storage: Store below 25°C. Protect from light.)

[FOR EXPORT ONLY]

Product Registration Holder (Malaysia) /

Manufactured by: HOVID Bhd.

121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar),
30010 Ipoh, Perak, Malaysia.

Revision date: December 2024

hovid