

HOVID - METHYL SALICYLATE LINIMENT

VIMET07-3var

DESCRIPTION

Clear, yellowish liquid.

COMPOSITION

Methyl Salicylate 25% v/v.

PHARMACODYNAMICS

Methyl Salicylate is used externally as a counterirritant-rubefacient. When it is rubbed on the skin, it produces mild irritation and is absorbed to an appreciable extent.

It is applied in liniments and ointments for the relief of pain in lumbago, sciatica and rheumatic conditions.

PHARMACOKINETICS

It is readily absorbed through the skin.

INDICATIONS

For relief of pain in lumbago, sciatica and rheumatic conditions.

WARNINGS AND PRECAUTIONS

This product contains methyl salicylate and when applied or rubbed on to the skin, can be absorbed through the skin into the blood. For patients taking warfarin, excessive application on to the skin for muscle or joint pains may increase the chances of bleeding.

It should not be applied to broken or inflamed skin or near to the eyes or mucous membranes.

MAIN SIDE/ADVERSE EFFECTS

Sensitivity reactions, urticaria and angioneurotic edema may occur in susceptible patients.

ROUTE OF ADMINISTRATION

Topical

CONTRAINDICATIONS

There are no data on contraindications in human.

SYMPTOMS AND TREATMENT OF OVERDOSE

Clinical features : Hyperventilation, hyperthermia, epigastric pain, nausea, vomiting, metabolic acidosis, pulmonary edema. Treat poisoning by emesis or gastric lavage, if appropriate. Electrolyte and metabolic abnormalities, if any, should be corrected.

DOSAGE AND ADMINISTRATION

Massage on intact skin 2 to 3 times a day as needed.

Note: The information given here is limited. For further information consult your doctor or pharmacist.

Storage:

Store below 30°C. Protect from light.

Presentation/Packing:

Liniment x 30ml, 60ml, 100 ml, 3.6 Litres.
(Not all presentations available locally)

Product Registration Holder / Manufactured by:

HOVID Bhd.,
121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar),
30010 Ipoh, Perak, Malaysia.

Revision date: July 2020