

70mm

**REXOM™ VITAMIN B-12 TABLET 50MCG****Name and Strength of Active Ingredients(s)**

Each tablet contains:

Cyanocobalamin ..... 50 mcg

**Product Description**

Pink colour, round with top score tablet.

**Pharmacodynamics**

This medicine contain cyanocobalamin vitamin B-12, which is used for the treatment of pernicious anaemia, and nutritional deficiencies of vitamin B-12 which results in macrocytic anaemia.

**Pharmacokinetics**Absorption

The absorption of cobalamins from the gut is dependent upon the glycoprotein intrinsic factor.

Distribution

Cobalamins are transported rapidly into the blood bound to protein, known as transcobalamins.

Elimination

Cobalamins are stored in the liver and excreted in the bile. They are known to cross the placenta.

**Indications**

Vitamin B-12 deficiency of dietary origin.

**Recommended Dose**

Adults and Elderly: One to three tablets daily.

Children: One tablet daily.

REXOM™ Vitamin B-12 tablet should be swallowed with water and taken between meals. Do not stop taking this medicine, except from doctor's advice. Your doctor may change the dose, depending on how you react to it.

**Route of Administration**

Oral.

**Contraindications**

Hypersensitivity to the product or any of the excipients.

**Warning and Precautions**

For pernicious anaemia, an adequate dose must be used and the blood picture must be examined regularly at least every three months for 18 months until stabilised, and then annually.

Indiscriminate administration of this medicine may mask precise diagnosis.

Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

**Interactions with Other Medicaments**

Absorption may be reduced by Para-aminosalicylic acid, colchicine, biguanides, neomycin, cholestyramine, potassium chloride, methyl dopa, and cimetidine.

Patients treated with chloramphenicol may respond poorly to this medicine.

Serum levels of this medicine may be lowered by oral contraceptives. These interactions are unlikely to have clinical significance.

Anti-metabolites and most antibiotics invalidate vitamins B-12 assays by microbiological techniques.

**Pregnancy & Lactation**

This medicine should not be used to treat megaloblastic anaemia of pregnancy because this is due to folate deficiency.

**Side Effects**

Sensitisation to this medicine is rare, but may present as an itching exanthema, and exceptionally as anaphylactic shock.

Acneiform and bullous eruptions have been reported rarely.

Patients who have become sensitised to this medicine by injection are often able to tolerate the oral route without trouble.

**Symptoms and Treatment of Overdose**

Overdosage is unlikely to require treatment.

**Effects on ability to drive and use machines**

None.

**Storage Condition**

Store below 30°C. Protect from heat and light. Keep container tightly closed.

**Dosage Forms and Packaging Available**

REXOM™ Vitamin B-12 Tablet 50mcg is packed in Alu-PVDC &amp; Alu-PVC blister pack.

Each blister are packed into 10 x 10's and 100 x 10's, in a carton along with a package insert.

REXOM™ Vitamin B-12 Tablet 50mcg is also packed in 90's and 100's per container. 100's is for export only.

**Registration Number:**

MAL19911374X

**Manufactured by & Product Registration Holder:****KCK Pharmaceutical Industries Sdn. Bhd.**

Plot 61, Bayan Lepas Industrial Park Phase 4,

Lintang Bayan Lepas 1, 11900 Bayan Lepas,

Pulau Pinang, Malaysia.

**Date of Revision**

7 January 2025

PI0073-2

160mm