

Axcel®
KAOPEC
SUSPENSION

COMPOSITION:

Each 5ml contains

Light Kaolin 1g
Pectin (Citrus) 20mg

Methylparaben 0.1%w/v as preservative.

PRESENTATION:

Cream coloured and cream (Butter) flavoured suspension.

INDICATION:

Antidiarrhoeal of non-specific origin.

DOSAGE & ADMINISTRATION:

For oral administration only.

Children 6 - 12 years : 20ml

To be taken 3 times a day, when necessary.

CONTRAINDICATION:

Contraindicated in intestinal obstruction.

PRECAUTIONS:

Appropriate fluid and electrolyte therapy should be given to protect against dehydration. Oral rehydration therapy - which is the use of appropriate fluids including oral rehydration salts - remains the most effective treatment for dehydration due to diarrhoea. The intake of as much as possible of these fluids is, therefore, imperative.

WARNING:

KAOLIN / PECTIN / KAOLIN-PECTIN IS NOT
RECOMMENDED FOR CHILDREN UNDER
6 YEARS OF AGE.

Severe constipation, which may lead to fecal impaction, may rarely occur in children and elderly patients taking kaolin and pectin. Kaolin and pectin may interfere with the absorption of other drugs, including antibiotics when administered concurrently.

SIDE EFFECT:

None.

STORAGE:

Keep container well closed. Store below 30°C.
Protect from light.

PACK QUANTITIES:

Available in 120ml amber glass bottle.

KEEP OUT OF REACH OF CHILDREN.

JAUHI DARI KANAK-KANAK.

Further information can be obtained from doctor, pharmacist or the manufacturer.

MAL19911416X

Product Registration Holder & Manufactured By:



Kotra Pharma (M) Sdn. Bhd.
No.1, 2 & 3, Jalan TTC 12, (90082-V)
Cheng Industrial Estate,
75250 Melaka, Malaysia.

Date of revision : 31/10/22