



DEXXON LOZENGES

Further information can be obtained from pharmacist, physician or the manufacturer.

COMPOSITION:

Each lozenge contains
Dequalinium Chloride 0.25mg

Sunset Yellow as colourant.

PRESENTATION:

Orange coloured and orange flavoured, round, flat lozenges, marked KOTRA logo on one side, 12.9mm in diameter.

INDICATIONS:

Minor infection of the mouth and throat.
Anginous infection : Angina catarrhalis, as adjuvant in angina lacunaris and Plaut-Vincent's angina.
Catarrhal infections : Pharyngitis, laryngitis.
Inflammation of the mucous membrane and gums : Gingivitis, ulceration stomatitis, aphthous stomatitis.

DOSAGE:

Adult and children : Allow 1 lozenge to be dissolved slowly in the mouth every 2 hours. After the inflammatory symptoms have subsided, suck 1 lozenge every 4 hours.

CONTRAINDICATION:

Patients hypersensitive to dequalinium chloride, patients being treated with sulphonamides and children under 3 years of age.

SIDE EFFECTS:

Side effects and undesirable accompanied symptoms have not been observed.

PRECAUTIONS:

Severe sore throat accompanied by high fever or headache, nausea and vomiting, any sore throat persisting more than 2 days may be serious, and need to be under medical attention.

STORAGE:

Keep container well closed. Store below 30°C. Protect from light.

KEEP OUT OF REACH OF CHILDREN

PACK QUANTITIES:

Available in blister pack of 100 x 10's, 2 x 10's & 10 x 10's.

MAL19911425X

Product Registration Holder & Manufactured By:



Kotra Pharma (M) Sdn. Bhd.
No.1, 2 & 3, Jalan TTC 12, (90082-V)
Cheng Industrial Estate,
75250 Melaka, Malaysia.

Date of Revision : 30/03/23