

Package insert

CHEWIES VITAMIN C 100MG CHEWABLE TABLET (MAL 19911658XZ)

Description	An orange coloured, orange flavoured, round and chewable tablet. 15mm in diameter and embossed with 'Chewies 100' on one side of the tablet.
Composition	Each tablet contains: Ascorbic Acid (Vitamin C) 100mg
Indication	For the prevention and treatment of Vitamin C deficiency.
Dosage	For oral administration. Children: 1-3 tablets daily to be sucked or chewed.
Pharmacokinetics	Ascorbic acid is readily absorbed from the gastrointestinal tract and is widely distributed in the body tissues. Plasma concentrations of ascorbic acid rise as the dose ingested is increased until a plateau is reached with doses of about 90 to 150mg daily. Body stored of ascorbic acid in health are about 1.5g although more may be stored at intakes above 200mg daily. The concentration is higher in leucocytes and platelets than in erythrocytes and plasma. In deficiency states the concentration in leucocytes declines later and at a slower rate, and has been considered to be a better criterion for the evaluation of deficiency than the concentration in plasma. Ascorbic acid is reversibly oxidised to dehydroascorbic acid; some are metabolised to ascorbate-2-sulfate, which is inactive, and oxalic acid which is the urine. Ascorbic acid in excess of the body's needs is also rapidly eliminated unchanged in the urine; this generally occurs with intakes exceeding 100mg daily. Ascorbic acid crosses the placenta and is distributed into breast milk. It is removed by haemodialysis.
Precautions	Use of ascorbic acid should be carefully considered when the following medical problem exists. Cystinuria, gout, hyperoxaluria or oxalosis, urate renal stones, Diabetes mellitus, glycose-6-phosphate dehydrogenase deficiency, hemochromatosis, sideroblastic anemia, thalassemia, sickle cell anemia
Drug interaction	Ascorbic acid may interfere with the effects of oral anticoagulants, anticholinergics and barbiturates
Contraindications	Unknown contraindications.
Side effects	Withdrawal scurvy may occur after prolonged administration of 2 to 3 grams per day. Occasionally, prolonged use of high doses of ascorbic acid may cause precipitation of urate, cysteine or oxalate, stones in the urinary tract.
Storage condition	Store in a dry place below 30°C. Keep bottle tightly closed. Keep out of reach of children. <i>Jauhi daripada kanak-kanak.</i>
Shelf life	4 years from date of manufacture.

Package insert

Product Registration Holder	AJ Research & Pharma Sdn Bhd (1014182-P) Unit 27.1, Level 27, Menara Felda,Platinum Park, No.11 Persiaran KLCC, 50088 Kuala Lumpur, Malaysia.
Manufactured by	Duopharma Manufacturing (Bangi) Sdn Bhd (53667-M) Lot 2 & 4, Jalan P/7, Section 13, Bangi Industrial Estate, 43650 Bandar Baru Bangi, Selangor. Malaysia
Revision date	October 2022

Further information can be obtained from your pharmacist, doctor or the manufacturer.