

HUACHI ENEMA

Patient Information Leaflet

Active Ingredients and Strength

Each 20ml contains :

Glycerin 6 gm

Sodium Chloride 3 gm

General Indication

The product is indicated for the relief of constipation

Contraindication

Hypersensitive to its active components

Side Effects/Adverse Reactions

The side effects tend to be minimal, but may include rectal discomfort, burning, and spasmodic bowel pain or stomach cramps. Some rectal bleeding or discharge also may occur

Precautions/Warnings

For rectal use only. Frequent or prolonged use of laxatives may result in dependence. Do not use when nausea, vomiting or abdominal pain is present. If use results in unusual pain or side effects, consult a physician. Keep this out of the reach of children. If sudden change in bowel habits persists over some time, consult a physician before using a laxative

Recommended Use

- 1) Remove the cap and press some liquid over the tip for easy insertion
- 2) Gently insert the tubular part into rectum completely
- 3) Squeeze applicator to empty all liquid into intestines
- 4) Remove applicator, cover with toilet tissue – remove toilet tissue when bowel movement become liquid
- 5) For better effectiveness, place applicator in warm water before using
- 6) Use 20ml for adults, 10ml for children

Pharmacology

Glycerin is readily absorbed from the intestine and is metabolised to carbon dioxide and glycogen or is used in the synthesis of body fats

Store below 30°C

Pack Size : 10ml, 20ml

Shelf Life : 5 years from date of manufacture

Sole Agent : Pembekalan Ubat Huathai Sdn Bhd

No. 17 Jalan Banang 3

Taman Johor

81200 Johor Bahru Johor, Malaysia