



WORMEX TABLET 125mg

Contents:

Wormex Tablet : Pyrantel Pamoate USP equivalent to Pyrantel base 125mg/tablet

Indications

For treatment of single or mixed infestations of pinworm/threadworm (Enterobius), roundworm (Ascaris), hookworm (Ancylostoma and Necator) and trichostrongyliasis.

Pharmacology

Pyrantel, a broad-spectrum anthelmintic, acts as a depolarising neuromuscular blocker, thereby causing sudden contraction, followed by paralysis of the helminths. The helminths are rendered unable to maintain their position in the intestinal lumen and are expelled from the body in the faecal stream by peristalsis.

Oral absorption is poor and incomplete and elimination is by renal and faecal excretions.

Adverse effects

Side effects occur only occasionally and are mild and transient; gastrointestinal disturbances, headache, dizziness, skin rash, and fever have been reported.

Precautions

- No special preparations (e.g. dietary restrictions or fasting, purging, or cleansing enemas) are required before, during, or immediately after therapy.
- Caution in patients with impaired liver function.
- Safety for use in pregnancy and lactation has not been established.

Drug interactions

Piperazine may antagonize the anthelmintic effects of pyrantel; concurrent use is not recommended.

Dosage and administration

Adults and children over 2 years:

Oral, 10mg (base) per kg of body weight with a maximum of 1 gram (base) as a single dose.

- For pinworm/threadworm infestation — Repeat in two to three weeks.
- For roundworm infestation — May be repeated in two to three weeks if required.
- For hookworm infestation — The dose may be given once a day for three days.

Note: 2.9 grams of pyrantel pamoate are equivalent to approximately 1 gram of base.

: Pyrantel may be taken on a full or empty stomach at any time of day.

Overdosage:

Because of poor gastrointestinal absorption, pyrantel serum concentrations are low. However, if overdose occurs, early gastric lavage and supportive therapy are necessary. There is no specific antidote.

Storage

: Keep container tightly closed.
Store in a cool, dry place, below 30°C, away from light.

Presentation/Packing

: Tablet 125mg 100's (EXPORT)

Manufactured & distributed by

: Prime Pharmaceutical Sdn. Bhd., Penang, Malaysia.