



PRI-CALCIUM TABLETS

Description: White, round and top-scored tablet.

Content

Calcium Lactate Pentahydrate 300 mg per tablet

Indications

As supplemental therapy for pregnant, lactating, and postmenopausal women; calcium deficiency and conditions requiring increased calcium intake.

Pharmacology

Calcium is essential to physiologic processes such as transmission of nerve impulses; contraction of cardiac, smooth, and skeletal muscles; renal function; respiration; blood coagulation; and other regulatory roles in the body.

Oral absorption decreases with age and increases by administration of vitamin D and it undergoes both renal and faecal excretions.

Adverse Effects

Constipation, gastrointestinal disturbances, and chalky taste may occur.

Precautions

- Caution in patients with impaired renal function or a history of renal stone formation.
- It should not be taken for prolonged periods without medical advice.

Contraindications

Avoid in patients with hypercalcemia, renal calculi, and ventricular fibrillation.

Drug Interactions

Calcium salts reduce the absorption of tetracyclines; Patient should be advised not to take calcium salts within 1 to 3 hour of tetracyclines.

Overdosage

Large doses may cause severe constipation, abdominal pain, muscle weakness, confusion and calcific renal stones.

Treatment consists of reducing or withdrawing calcium intake and administering large amounts of fluid to combat dehydration and reduce renal deposition of calcium; or as directed by physician.

Dosage and Administration

Adults : Oral, 300 to 600 mg three times daily with meals, or as required.

Storage : Keep container tightly closed. Store in a cool, dry place, below 30°C, away from light.

Presentation/Packing : Tablet 300 mg x 150's

Shelflife : 3 years

Manufactured & : **PRIME PHARMACEUTICAL SDN.BHD.**

Distributed by 1505, Lorong Perusahaan Utama 1,
Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia.