

A.N.B. Dextrose 10% in Dis. Water Inj

Controlled Medicine/*Ubat Terkawal*

Each 100 mL contains

Dextrose	10 g
Sterile Water for Injection ad to	100 mL

Pharmacodynamics/Pharmacokinetics Time to peak effect: 8 to 12 hours.

Indication Fluid and nutrient replenisher.

Adverse effects

It may cause febrile response; infection at the site of injection; tissue necrosis; venous thrombosis or phlebitis extending from the site of injection; extravasation; hypovolemia; hypervolemia; dehydration; mental confusion or unconsciousness. These may occur because of the solution or administration technique. Use the largest available peripheral vein and a well placed small bore needle.

Precautions

Perform clinical evaluations and laboratory determinations to monitor fluid balance, electrolyte concentrations and acid-base balance.

Vitamin B complex deficiency may occur with dextrose administration.

Warnings

Dextrose solution IV can cause fluid or solute overload resulting in dilution of serum electrolyte concentrations, overhydration, congested states or pulmonary edema. Use dextrose-containing solutions with caution in patients with subclinical or overt diabetes mellitus or carbohydrate intolerance.

Contra-indication

It is contraindicated in diabetic coma while blood sugar is excessively high. Hypertonic dextrose solutions are contraindicated in patients with anuria, or intraspinal or intracranial hemorrhage, and in patients with delirium tremens if such patients are already dehydrated. Hypertonic dextrose solutions are also contraindicated in patients with diabetic coma or known allergy to corn or corn products.

Drug interactions

Cautiously administer parenteral fluids, especially those containing sodium ions, to patients receiving corticosteroids or corticotropin.

Recommended Dosage Intravenous infusion or as physician's order.

Treatment of Overdosage

In the event of a fluid or solute overload during parenteral therapy, reevaluate the patient's condition and institute appropriate corrective treatment.

Storage conditions

Store below 30°C.

Keep out of reach of children

"Discard unused portion"

Packaging

Polypropylene bottle of 20 x 500 mL

Description of product Sterile, clear, colourless solution.

Name and address of actual manufacturer

A.N.B. Laboratories Co., Ltd.
557 Ramintra Rd., Bangkok, Thailand.

Product Registration Holder

Delfi Marketing Sdn. Bhd. (15872-P)
Level 6, Block A, Sky Park One City,
Jalan USJ 25/1, 47650 Subang Jaya,
Selangor, Malaysia.

Date of Revision

17/06/2019