

อ้างอิง ใบคำเนนการจัดทำวัสดุบรรจุหีบห่อใหม่หรือแก้ไขวัสดุบรรจุหีบห่อเดิม เลขที่ 026/63		ตรวจครั้งที่	1
ชื่อแบบ	เอกสารกำกับยา A.N.B. Dextrose 5% in NS 0.9% Inj (Malaysia)	กระดาษ	
พิมพ์ (สี)	1 สี (ดำ)	ขนาด	95x130 mm
หมายเหตุ		รหัสวัสดุ	001546-D02
ผู้จัดทำ	สุนันทา กุลไพศาล	ผู้ตรวจแบบ (Approved by)	
วันที่จัดทำ	27/02/63	4. จัดซื้อ วันที่/...../..... 5. ผลิต วันที่/...../..... 6. QC วันที่/...../..... 7. QA วันที่/...../.....	
หน้า	1/1	1. การตลาด วันที่/...../..... 2. ต่างประเทศ..... วันที่/...../..... 3. ทะเบียน วันที่/...../.....	

A.N.B. Dextrose 5% in NS 0.9% Inj

Controlled Medicine/Ubat Terkawal

Each 100 mL contains:

Dextrose

5 g

Sodium Chloride

0.9 g

Sterile Water for Injection

q.s. to 100 mL

Product Description:

Sterile, clear, colourless solution

Pharmacodynamics/Pharmacokinetics:

Time to peak effect: 8 to 12 hours.

Indication:

Fluid, nutrient and electrolyte replenisher.

Recommended Dosage:

Intravenous infusion or as physician's order.

Contraindications:

It is contraindicated in diabetic coma while blood sugar is excessively high. Hypertonic dextrose solutions are contraindicated in patients with anuria, or intraspinal or intracranial haemorrhage, and in patients with delirium tremens if such patients are already dehydrated. Hypertonic dextrose solutions are also contraindicated in patients with diabetic coma or known allergy to corn or corn products.

Warnings:

Dextrose solution IV can cause fluid or solute overload resulting in dilution of serum electrolyte concentrations, overhydration, congested states or pulmonary edema. Use dextrose-containing solutions with caution in patients with subclinical or overt diabetes mellitus or carbohydrate intolerance.

Precautions:

Perform clinical evaluations and laboratory determinations to monitor fluid balance, electrolyte concentrations and acid-base balance. Vitamin B complex deficiency may occur with dextrose administration.

Drug Interactions:

Cautiously administer parenteral fluids, especially those containing sodium ions, to patients receiving corticosteroids or corticotrophin.

Adverse Effects:

It may cause febrile response; infection at the site of injection; tissue necrosis; venous thrombosis or phlebitis extending from the site of injection; extravasation; hypovolemia; hypervolemia; dehydration; mental confusion or unconsciousness. These may occur because of the solution or administration technique. Use the largest available peripheral vein and a well-placed small bore needle.

Treatment of Overdose:

In the event of a fluid or solute overload during parenteral therapy, re-evaluate the patient's condition and institute appropriate corrective treatment.

Storage Conditions:

Store below 30°C. Keep out of reach of children. Discard unused portion.

Packaging:

- Polypropylene bottle of 20 x 500 mL

- Polypropylene bottle of 10 x 1000 mL

Manufacturer:

A.N.B. Laboratories Co., Ltd.

557 Ramintra Rd., Bangkok, Thailand.

Product Registration Holder:

Delfi Marketing Sdn. Bhd.

Level 6, Block A, Sky Park One City,

Jalan USJ 25/1, 47650 Subang Jaya,

Selangor, Malaysia.

Date of Revision:

8 April 2019

Page 1 of 1

001546-D02