

Hyalgan® 20mg/2ml injection

(Sodium Hyaluronate)
For intra-articular use

COMPOSITION - Highly viscous aqueous solution of a defined molecular fraction of purified hyaluronic acid, 10 mg/ml, for intra-articular administration.

Each syringe contains:

- Hyaluronic acid sodium salt (Hyalectin®)	20.0 mg
- Sodium chloride	17.0 mg
- Monobasic sodium phosphate 2H ₂ O	0.1mg
- Dibasic sodium phosphate 12H ₂ O	1.2 mg
- Water for injection q.s. to	2.0 ml

DESCRIPTION - Hyaluronic Acid, a natural polymer of the glycosaminoglycan family, is an important component of the extracellular matrix and is present in a particularly high concentration in cartilage and synovial fluid. The active substance in HYALGAN® is a defined fraction of sodium hyaluronate, obtained by a molecular filtration procedure from biological material.

PHARMACOLOGY - The intra-articular administration of HYALGAN® into arthritic joints with degenerating cartilage surfaces and pathologically altered synovial fluid improves joint function, due to the normalization of synovial fluid quality and the activation of tissue repair processes in articular cartilage.

TOXICOLOGY - The LD₅₀ was greater than 100 mg/kg in the rat and mouse. Results of subacute, chronic and foetal toxicity studies, fertility studies, peri- and post-natal toxicity, mutagenesis, immunogenicity and intra-articular tolerability studies showed that the product was well tolerated.

PHARMACOKINETICS - Hyaluronic acid is efficiently metabolized in the animal organism. The substance is rapidly removed from the blood stream and quickly degraded in the liver. This finding is in accordance with the biological nature of the substance.

After intra-articular administration of labelled hyaluronic acid in laboratory animals, maximum blood levels of radioactivity are seen after about 48 hours. The radioactivity is excreted mainly in the urine. About 42% of the administered dose is present in the liver 24 hours after administration. Radioactivity is seen in the synovia within 2 hours and in the articular cartilage within 6 hours. The cartilage is the most stable location. In the synovial fluid the substance persists for 4 to 5 days.

INDICATIONS - Traumatic and degenerative joint disease. - Adjuvant in orthopaedic surgery.

CONTRA-INDICATIONS - Individual hypersensitivity to the product. To date, no other contra-indications to the intra-articular administration of hyaluronic acid are known.

ADVERSE EXPERIENCE - HYALGAN® is generally well tolerated. However in some patients swelling of the joint and/or local pain may occur after intra-articular injection. Such symptoms are transient and will disappear after a few hours.

DOSAGE AND ADMINISTRATION - According to the size of the joint, 2 ml of HYALGAN® (20 mg) or less administered intra-articularly once a week for 3 to 5 weeks, depending on the severity of the disorder. More than one joint may be treated at the same time.

WARNINGS AND PRECAUTIONS - It is necessary to follow a correct technique of intra-articular injection in accurately aseptic conditions, taking care not to cause tissue damage in the injection path.

USE DURING PREGNANCY AND LACTATION - To date, no contra-indications to the use of the product in pregnancy and during lactation are known.

OVERDOSAGE - Phenomena due to overdosage have not been noted to date.

DRUG INTERACTIONS - No interaction with other drugs have been noted to date.

SUPPLY - One 2 ml pre-filled syringe.

STABILITY AND STORAGE RECOMMENDATIONS

This product has a shelf life of 36 months if stored in its original package, protected from light below 30°C. Do not freeze.

KEEP OUT OF REACH OF CHILDREN

Date of revision: 21/10/2020



MANUFACTURED BY:

Fidia Farmaceutici S.p.A., Abano Terme, Italy.