

# SCMC

## PLUS SYRUP

### 250mg/5ml



**SCMC PLUS** Syrup 250mg/5ml contains S-carboxy-methylcysteine, an active mucolytic agent. It is formulated for adult with wet cough where there is excessive phlegm present. It loosens phlegm so it can be coughed out easily.

| Contents                 | Strength  | Preservatives                                     |
|--------------------------|-----------|---------------------------------------------------|
| S-carboxy-methylcysteine | 250mg/5ml | Methylparaben 0.21%w/v<br>Propylparaben 0.031%w/v |

#### PRODUCT DESCRIPTION:

A yellow, fruity flavoured homogeneous syrup.

#### PHARMACODYNAMICS:

S-carboxymethylcysteine is an active mucolytic agent. It dissolves mucus plugs, relieves tracheo-bronchial congestion and improves respiration thus facilitating expectoration.

#### PHARMACOKINETICS:

S-carboxymethylcysteine is absorbed from the gastro-intestinal tract excreted in the urine as unchanged drug and metabolites.

#### INDICATION:

Reduction of sputum viscosity to facilitate expectoration and evacuation.

#### DOSAGE:

Adult and children above 12 years: 15ml three times daily.

#### SIDE EFFECTS:

Nausea, headache, gastric discomfort, diarrhoea, gastro-intestinal bleeding, and skin rash have occasionally occurred with S-carboxymethylcysteine.

#### Immune System Disorders

Anaphylactic / anaphylactoid reaction.

#### Skin And Subcutaneous Tissue Disorders

Severe cutaneous adverse reactions (SCAR) e.g. erythema multiforme, Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN). In most of these cases reported at least one other drug was administered at the same time, which may have possibly enhanced the described mucocutaneous effects.

#### WARNINGS AND PRECAUTIONS:

It is recommended that S-carboxymethylcysteine be used with caution in patients with a history of peptic ulcer and be avoided in patients with active ulceration.

**SCMC Syrup** may cause severe allergy and serious skin reactions. Stop using **SCMC Syrup** and seek medical assistance immediately if you experience any of the following symptoms:

- Severe allergy: breathing difficulties, light headedness, skin swellings or rash.
- Severe skin reaction: skin reddening, blisters, rash, fever, sore throat or eye irritation.
- Do not use for more than 5 days.
- If cough persists, please consult your doctor.

#### CONTRAINDICATIONS:

Contraindicated in children below 2 years of age.

#### PACKING:

100ml

#### STORAGE CONDITION:

Keep container tightly closed.  
Store in a dry place below 30°C.  
Protect from light.  
Keep out of reach of children.  
Jauhi daripada kanak-kanak.

Shelf life: Please refer to outer package.

Route of Administration: Oral

Product Registration Holder :

**DUOPHARMA MANUFACTURING (BANGI) SDN. BHD.**

Lot No 2, 4, 6, 8 & 10, Jalan P/7, Section 13,  
Bangi Industrial Estate, 43650 Bandar Baru Bangi,  
Selangor, Malaysia.

Manufacturer :

**DUOPHARMA MANUFACTURING (BANGI) SDN. BHD.**

Lot No 2 & 4, Jalan P/7, Section 13, Bangi Industrial  
Estate, 43650 Bandar Baru Bangi, Selangor, Malaysia.

Revision Date: Oct 2025  
1500010203 N.2

Size: 148mm (w) x 210mm(h)



Black 100%