

KAOLIN-PEC SUSPENSION
(Antidiarrhoeal)**Active Ingredients:**

Each 30ml contains -
Light Kaolin 6g
Pectin 150mg
Preservatives -
Methyl Hydroxybenzoate 0.1%w/v
Propyl Hydroxybenzoate 0.01%w/v
Sodium Benzoate 0.2%w/v

Presentation:

Bottle of 60ml.

Product Descriptions:

Plum flavoured, creamy brown suspension.

Pharmacology:

Light Kaolin together with Pectin is of value in the symptomatic treatment of diarrhoea caused by agents capable of being adsorbed e.g. in the diarrhoea of food poisoning or dysentery. These agents are adsorbed from the alimentary tract by Light Kaolin and the bulk of the faeces are increased.

Indications:

Control of diarrhoea by adsorbing toxic substances and increasing bulk of faeces.

Dosage and Administration:

Adults and children above 12 years:

Take 3 tablespoons (30ml) 3 times a day or when needed.

Children 6 to 12 years:

Take 4 teaspoons (20ml) 3 times a day or when needed.

Shake well before use.

Warnings:

Kaolin/Pectin/Kaolin-Pectin is not recommended for children under 6 years of age.

This preparation contains Sodium Metabisulphite that may cause serious allergic type reactions in certain susceptible patients. Do not use if known to be hypersensitive to bisulphites.

Severe constipation, which may lead to fecal impaction, may rarely occur in children and the elderly patients taking Kaolin and Pectin.

Precautions:

Appropriate fluid and electrolyte therapy should be given to protect against dehydration. Oral rehydration therapy, which is the use of appropriate fluids combined with oral rehydration salts, remains the most effective treatment for dehydration due to diarrhoea.

The intake of as much of these fluids as possible is therefore imperative.

Interactions with Other Medicaments:

The absorption of Lincomycin has been reported to be significantly reduced by kaolin.

Kaolin and pectin may interfere with the absorption of other drug, including antibiotics, administered concurrently.

Undesirable Effects:

The adsorbent properties of kaolin may influence the gastro-intestinal absorption of other drugs.

Storage:

Store below 30°C. Protect from light. Keep cap tightly closed.

Keep out of reach of children. Jauhkan daripada capaian kanak-kanak.

Shelf Life:

3 years from the date of manufacture.

Manufacturer and Product Registration Holder

WINWA MEDICAL SDN. BHD.

198101009677(75794-A)

1952, Lorong I.K.S. Bukit Minyak 3,

Taman I.K.S. Bukit Minyak,

14000 Bukit Mertajam,

Pulau Pinang, Malaysia.

MAL19912826X

Date of Revision: 28 August 2025

X.LKL-1.08

Size: 75mm x 100mm



Dark Blue (Pantone 2767U)

Checked by	
Approved by	