

WARI ETHYLCHLORIDE SPRAY

General characteristics: The spray tin contains ethylchloride, a clear, colourless, liquefied gas.

WARI ETHYLCHLORIDE SPRAY is applied in local skin anesthesia for a great number of indications. Its anesthetic effect is achieved by the spraying process, because ethylchloride, previously liquefied by compression, evaporates and consequently a coldness is produced which leads to a refrigeration and brings about a numbness of the sprayed areas of the skin and thus a pain alleviation.

WARI ETHYLCHLORIDE SPRAY is supplied in handy and attractive spray tins which offer advantages in comparison with ampoules or glass bottles (unbreakable, flat bottom). The spray tin does not contain any propellant. The spraying head guarantees a point-shaped stream by which an optimal application is achieved.

In local skin anesthesia WARI ETHYLCHLORIDE SPRAY is sprayed onto the areas of the skin to be treated from a distance of about 30 cm until a fine white film is being produced - thereafter do not continue spraying to avoid skin burning.

MEDICAL INFORMATION - PLEASE READ CAREFULLY!

Composition: 100 ml liquefied gas correspond to 88 g ethylchloride.

Indications: Local skin anesthesia.

Contra-Indications: Unknown.

Precaution: WARI ETHYLCHLORIDE SPRAY must not be used for short-term- or full narcoses, as it may cause severe damage to heart, liver and kidneys. Do not spray onto open injuries or wounds!

Side effects: After application of this spray there might occur allergic skin reactions.

Interactions: Unknown.

Dosage and mode of application: Use outdoors or in well ventilated rooms only. WARI ETHYLCHLORIDE SPRAY is only for external use in local skin anesthesia. If not otherwise prescribed, WARI ETHYLCHLORIDE SPRAY is sprayed onto the areas of the skin to be treated from a distance of approx. 30 cm until a fine white film is being produced.

Hint: In case of application in the region of head or neck special precautions have to be taken in order to avoid that ethylchloride is inhaled. **Duration of use:** Mostly one single application of WARI ETHYLCHLORIDE SPRAY is sufficient. The physician has to decide whether the treatment is to be repeated. After the expiry date WARI ETHYLCHLORIDE SPRAY should not be used any longer.

Storage: To be stored at 15° to 25°C out of the reach of children and protected from sunlight. Do not use near open fire and do not spray it on glowing objects. Do not force open or burn the tin.

Presentation: Spray tin of 100 ml.



WALTER RITTER - GmbH + Co. KG
Germany