

Package Insert

1. Product name

Senna 7.5 mg tablets

2. Name and Strength of Active Ingredient

Each tablet contains 7.5 mg sennosides.

Excipient(s) with known effect

This product contains 121.8 mg lactose.

3. Product description

Tablet.

Brown, round, scored tablets with Remedica's logo on one side. The shade of the colour may vary from batch to batch.

The score line is not intended for breaking the tablet.

4. Pharmacological Properties

Pharmacodynamics

Pharmacotherapeutic group: Drugs for constipation, ATC code: A06AB06

The sugar moiety of the sennosides is removed by bacteria in the large intestine releasing the active anthrone fraction. This stimulates peristalsis via the submucosal and myenteric nerve plexuses. Senna acts in 8-12 hours.

Pharmacokinetics

The action of the sennosides is colon specific and does not depend upon systemic absorption.

5. Clinical Particulars

Indications

Treatment of occasional constipation.

Posology and method of administration

Posology

Adults only: 2-4 tablets in 24 hours, to be taken at night.

Paediatric population

Not to be given to children except on medical advice.

Method of administration

Oral administration.

Contraindications

Senna is contra-indicated in patients with hypersensitivity to the active substance or to any of the excipients listed in section 2.0. Senna should not be used when symptoms of appendicitis, intestinal obstruction, inflammatory bowel disease, including ulcerative colitis, or abdominal pain of unknown origin are present.

Warnings and Precautions

If symptoms persist, consult your doctor.

Senna should not be used for prolonged periods.

Senna 7.5 mg tablets contain lactose. Patients with rare hereditary problems of galactose intolerance, the Lapp lactose deficiency or glucose-galactose malabsorption should not take this medicine.

Interactions with other medicaments

None known.

Pregnancy and Lactation

Pregnancy

The product should not be used during pregnancy and lactation unless considered essential by the physician.

Breast-feeding

Anthraquinones are excreted into the milk of nursing mothers but clinical studies have shown that breast-fed infants of mothers taking sennosides in the puerperium did not show any side effects to the drug.

Undesirable effects

Senna may give rise to griping abdominal pain.

Symptoms and treatment of overdose

Where diarrhoea is severe conservative measures are usually sufficient; generous amounts of fluid, especially fruit drinks, should be given.

6. Pharmaceutical Particulars

Storage conditions

Do not store above 30°C. Protect from light and moisture.

Shelf-life

5 years.

Pack size available

This product is available in pack sizes of: 60 (6 x 10) and 1000 (100 x 10) tablets in blister packs.

7. Manufacturer

Remedica Ltd.
Aharnon Street, Limassol Industrial Estate,
3056 Limassol, Cyprus

8. Product registration holder

Eucogen Sdn Bhd.,
No. 6A, Jalan Sungai Burung U 32/U, Bukit Rimau, Seksyen 32, 40460 Shah Alam.

NOVA Borneo Company Sdn Bhd,
Unit 4GF, Bangunan Zainuddin & Azizah,
Spg 501, Jalan Tutong, Kg Telanai, Brunei-Muara BA2312, Brunei Darussalam

9. Date of revision of the text

15/05/2025

For internal use only: my-pi-senna-tabs-a2