

WELLMEX HYDROGEN PEROXIDE 6% w/v SOLUTION

Description	: A colourless odorless liquid
Composition	: Hydrogen Peroxide 6%
Indications	: As first aid in the treatment of cuts, fresh wounds and abrasions. To remove dressings from wounds. As a gargle or mouthwash.
Direction for Use	: First aid – In the treatment of cuts, fresh wounds and abrasions, dilute the peroxide with an equal volume of water and apply on lint or cotton wool to arrest bleeding and disinfect the tissue. Allow the oxygen to escape and lightly fix the dressing in position with a bandage or strip of adhesive plaster. To remove dressings from wounds- Dilute the peroxide with an equal volume of water. Thoroughly soak the dressing with this solution and leave for some minutes before removal. For gargle or mouthwash- Use one dessert spoonful of peroxide in half a tumblerful of water as necessary. Catarrh – Mix one teaspoon of peroxide with a quarter of a tumblerful of water and sniff up the nose.
Route of Administration	: For topical use only
Pharmacodynamic properties	: Hydrogen peroxide is used as a disinfectant. It releases oxygen when applied to tissues. The effect lasts only as long as the oxygen is being released and is of short duration. The anti-microbial effect of the liberated oxygen is reduced in the presence of organic matter. It is used to cleanse wounds and ulcers in concentrations of up to 6%. Adhering and blood soaked dressings may be released by the application of a solution of hydrogen peroxide.
Use in pregnancy and lactation	: Hydrogen Peroxide should be avoided during pregnancy and lactation. No safety data is available for these conditions.
Contraindication	: Not known
Precautions / Warnings	: It is dangerous to inject hydrogen peroxide into closed body cavities from which the released oxygen has no free exit.
Interaction with Other Medicaments	: Not known
Side Effects/Adverse Reaction	: Strong solutions of hydrogen peroxide produce irritating burns on the mucous membrane but the pain disappears in about an hour.
Symptoms and Treatment for Overdose	: Not known

Pack Size	: 30 ml, 60 ml and 450 ml
Storage Conditions	: Store below 30 °C and away from direct heat and light. Keep out of reach of children. Jauhi daripada kanak-kanak
Shelf Life	: 3 years
Name and Address of Manufacturer	: The Government Pharmaceutical Organization, 75/1, Rama 6 Road, Ratchathewi, Bangkok, 10400 Thailand.
Product Registration Holder	: NGL TRADING SDN. BHD., (1020904-A) 4-1-5, Kompleks Kantonmen Prima, Batu 4 ½, Jalan Ipoh, 51200 Kuala Lumpur, Wilayah Persekutuan Kuala Lumpur.
M’sia Reg. No.	: MAL19913109X
Revision Date	: 06.11.2025

THE GOVERNMENT PHARMACEUTICAL ORGANIZATION
75/1 Rama VI Rd., Ratchathewi, Bangkok, Thailand