For the use only of a Registered Medical Practitioner or a Hospital or a Laboratory.

ELECTRAL FORTE

GRANULES

COMPOSITION

Electral Forte Granules 6g:
Each sachet of 6 g contains
Sodium Chloride BP .....0.7 gm
Potassium Chloride BP ..... 0.3 gm
Sodium Citrate BP .....0.58 gm
Anhydrous Glucose BP .....4.0 gm
Sodium Benzoate.....0.004 gm
(preservative)
Excipients .....q.s.

6 g of Electral forte in 200 ml of solution supplies:
Sodium.....90 mmols/L
Potassium.....20 mmols/L
Chloride..... 80 mmols/L
Citrate.....10 mmols/L
Glucose.....111 mmols/L
Total.....311 mmols/L

Electral Forte Granules 30g
Each sachet of 30 g contains
Sodium Chloride BP.....3.5 gm
Potassium Chloride BP ..... 1.5 gm
Sodium Citrate BP ..... 2.9 gm
Anhydrous Glucose BP..... 20.0 gm
Sodium Benzoate.....0.02 gm
(preservative)
Excipients ..... q.s.

30 g of Electral forte in 1000 ml of solution supplies:
Sodium.....90 mmols/L
Potassium.....20 mmols/L
Chloride..... 80 mmols/L
Citrate.....10 mmols/L
Glucose.....111 mmols/L
Total.....311 mmols/L

PHARMACEUTICAL FORM
Powder for reconstitution

DESCRIPTION
Off white granular powder having orange like aroma, solution in water gives sweet, sour taste.

PHARMCOLOGICAL DATA
Mechanism of Action:
Electral forte is an Oral Rehydration Salts (O.R.S) Solution. The key constituents of ORS are sodium and glucose. The central principle of oral rehydration therapy (ORT) is the utilization of sodium-glucose co-transport in the small intestine, a phenomenon which remains largely unaffected during acute infectious diarrhea. Thus the success of ORT is largely dependent on glucose-driven sodium absorption (Transcellular route) leading to passive absorption of water by the paracellular route. The clinical result is usually rapid rehydration and correction of acidosis.

Pharmacokinetic & Bioavailability
No information is available.

INDICATIONS
It is indicated for the oral correction of fluid and electrolyte loss in infants, children and adults. Treatment of watery diarrhoea of varying aetiologies including gastroenteritis in all age groups.

CONTRAINDICATIONS
The preparation is contraindicated in patients with dehydration of 10% or more and continued vomiting.

DOSAGE & ADMINISTRATION
Oral rehydration solutions are used for oral replacement of electrolytes and fluids in patients with dehydration, particularly that associated with acute diarrhea of various aetiologies.
The dosage of oral rehydration solutions should be tailored to the individual based on body-weight and the stage and severity of the condition. The initial aim of treatment is to rehydrate the patient, and, subsequently, to maintain hydration by replacing any further losses due to continuing diarrhea and vomiting and normal losses from respiration, sweating, and urination.
For adults 200 to 400 ml of oral rehydration solution for every loose motion has been suggested. The dosage for children is 200ml for every loose motion, and for infants is 1 to 1.5 times their usual feed volume. Normal feeding can continue after the initial fluid deficit has been corrected. Breast feeding should continue between administrations of oral rehydration solution.

Dilution
6g of Electral forte is to be dissolved in 200ml of previously boiled and cooled water.
30g of Electral forte is to be dissolved in one liter of previously boiled and cooled water.

SPECIAL POPULATIONS
Pregnancy and Lactation:
There are no special problems with pregnant & lactating women.
Neonates:
Although any moderately dehydrated individual is at risk of adverse effects, particularly if ORS are used inappropriately, there should be special concern about this group.
Children:
As for neonates, there should be special concern for young infants, but there are no special problems with other children.
The elderly:
There are no special problems with this group.
Concurrent disease:
There should be special concern about patients who are hypernatremic before treatment.

PRECAUTIONS & WARNINGS
Precautions:
Oral rehydration salts should be reconstituted only with water and at the volume stated. Fresh drinking water is generally appropriate, but freshly boiled and cooled water is preferred when the solution is for infants. The solution should not be boiled after it is prepared. Other ingredients such as sugar should not be added. Unused solution should be stored in a refrigerator and discarded 24 hours after preparation.
Warnings:
Oral rehydration solutions are not appropriate for patients with gastro-intestinal obstruction, oliguric or anuric renal failure or when parenteral rehydration therapy is indicated as in severe dehydration or intractable vomiting.

DRUG INTERACTIONS
No drug interactions have been described for oral rehydration solution.

SIDE EFFECTS / ADVERSE REACTIONS
Vomiting can occur after administration of oral rehydration solution, and may be an indication that it was administered too quickly. If vomiting occurs, administration should be halted for 10 minutes then resumed in smaller, more frequent, amounts.
The risk of hypernatraemia or overhydration after administration of oral rehydration solutions is low in patients with normal renal function. Overdosage of oral rehydration solutions in patients with renal impairment may lead to hypernatraemia and hyperkalaemia.

OVERDOSAGE & TREATMENT OF OVERDOSAGE
No information is available on overdosage & treatment of overdosage with oral rehydration solution.

TOXICOLOGY
The main constituents of oral rehydration salts are physiological and when used appropriately are not toxic.

STORAGE CONDITIONS
Store below 30°C.

SHELF LIFE
36 months from date of manufacture.

PRESENTATION
For Electral Forte Granules 6g and 30g: 20 sachets of 6g or 30g granules are packed in a printed carton along with a package insert.

Revision Date: 25th June 2025

MAL No.: MAL19913278XZ

Manufactured in India by:
FDC Limited
B-8, MIDC Industrial Area, Waluj,
Chhatrapati Sambhajinagar - 431 136,
Maharashtra.

Marketing Authorization Holder/Imported by:
UNIMED SDN BHD
53, Jalan Tembaga SD 5/2B,
Bandar Sri Damansara,
52200, Kuala Lumpur, Malaysia.

2000016624

FDC Limited	LEAFLET	Mini Pharma Code no.: **Front: 1772 / Back: 1771**
Size	110 (L) x 144 (H) mm	
Folded Size	----------	
No. of Folds	----------	
GSM	60 gsm	
Paper	Maplitho	
Specification No.	----------	
Colour	■ Pantone Black C	
SAP Code: 2000016624	Supercedes SAP Code : 2000006630	
Location: Waluj - Malaysia	Prepared On: 25-06-2025	

Reason for Change:
1) Change in Indications.
2) Preservatives added.
3) Aurangabad word replace with Chhatrapati Sambhajinagar.
4) Date of revision updated.